

March 1833.

Part 191,

*Price Two Shillings and Sixpence,*

TO BE CONTINUED MONTHLY,

OF THE

**Botanical Cabinet,**

CONSISTING OF

COLOURED DELINEATIONS

OF

PLANTS

FROM ALL COUNTRIES:

WITH

A SHORT ACCOUNT OF EACH, DIRECTIONS FOR  
MANAGEMENT, &c.

BY

CONRAD LODDIGES & SONS.

—  
THE PLATES BY  
GEORGE COOKE.  
—

*Each Part will contain Ten Plates, and Ten  
Parts will form one Volume.*

London:—Published by  
JOHN AND ARTHUR ARCH, CORNHILL;  
LONGMAN, REES, ORME, BROWN,  
AND CO., PATERNOSTER-ROW;  
AND C. LODDIGES AND SONS, HACKNEY.

—  
1833.  
—



Rape Book  
S B 407  
1367  
1833  
V. 20

THE  
BOTANICAL CABINET

Consisting of  
Coloured Delineations

OF  
Plants  
from all Countries,

with a short Account of each.  
Directions for Management &c. &c.

CONRAD LODDIGES & SONS

PL. XX.

The Plates by

GEORGE COOKE.

"Even Solomon in all his glory  
was not arrayed like one of these."


London, Published by John & Arthur Arch, Cornhill:  
Longman, Rees, Orme, Brown & Green, Paternoster Row:

and C. Loddiges & Sons, Hackney,  
1833.

NO. BOT. GARDEN

1911





*Euphorbia punicea.*



No. 1901.

**EUPHORBIA PUNICEA.**

Class.

**DODECANDRIA.**

Order.

**TRIGYNIA.**

.....

This is a native of Jamaica, whence it was introduced in 1778. It is a handsome, erect, shrubby species, growing sometimes to the height of six or seven feet. It flowers freely, and at various seasons, when its appearance is very splendid.

With us it is necessary to preserve it constantly in the stove. It may be increased, although with difficulty, by cuttings, and sometimes bears seeds. The soil should be light loam.





*Hedychium stenopetalum*



No. 1902.

**HEDYCHIUM STENOPETALUM.**

Class.	Order.
<i>MONANDRIA</i>	<i>MONOGYNIA.</i>

.....

This was introduced lately by Dr. Wallich, from India. It grows to the height of six feet or more, having leaves about a foot and a half long, and flowered with us in the month of March, 1832.

Like the other species, it requires the stove, and to be cultivated in perfection should have a large sized pot and rich loamy soil. It will increase by separating its knobby roots early in the spring.





*W. L. C. del.*

*Sparaxis tricolor.*



No. 1903.

**SPARAXIS TRICOLOR.**

Class.	Order.
<b>TRIANDRIA</b>	<b>MONOGYNIA.</b>

.....

This is a native of the Cape of Good Hope, and was introduced in 1789. Its flowers are most splendid in their vividly contrasted colours, and it is a great ornament of the greenhouse when in perfection; or it may be kept very well in a border close to the front of a stove, with no other protection. It increases by offsets from the bulbs, or by seeds, which are often produced. The soil should be sandy peat.





*Maxillaria ochroleuca*



No. 1904.

**MAXILLARIA OCHROLEUCA.**

Class.	Order.
<i>GYNANDRIA</i>	<i>MONANDRIA.</i>

... ..

A native of Brazil, lately introduced : it grows vigorously, and flowers during summer and autumn. It is not so splendid as some of this genus, yet well deserves cultivation, which is not difficult planted in moss and vegetable earth, with fragments of pot intermixed.

It will occasionally admit of separation for increase, and must be always kept in the stove.





*Hodder & Goulet del.*

*Blechnum gracile.*



No. 1905.

**BLECHNUM GRACILE.**

Class.	Order.
<i>CRYPTOGAMIA</i>	<i>FILICES.</i>

.....

This pretty little fern is a native of Brazil : we received it in 1830 from our friend, Mr. Otto, of Berlin, and have kept it in the stove, in which it appears to thrive. The drawing was made in December.

It should be potted in sandy peat, and will occasionally increase by dividing the roots.





*Chytra nobilis*



No. 1906.

**CLIVIA NOBILIS.**

Class.	Order.
<i>HEXANDRIA</i>	<i>MONOGYNIA.</i>

.....

This splendid plant is a native of the Cape of Good Hope, and was discovered and introduced by Mr. Bowie, as we are informed by Mr. Lindley, who has given an elegant figure of it in the Botanical Register, and named it in compliment to her Grace the Duchess of Northumberland, in whose princely garden at Syon it first flowered.

Our plant was in perfection in the months of June and July, and appears likely to ripen its seeds, by which of course it may be multiplied: it does also occasionally produce offsets. We have hitherto kept it in the stove, potted in sandy peat earth, in which it thrives very well.







No. 1907.

**MENZIESIA POLIFOLIA** *nana*.


Class.	Order.
<b>OCTANDRIA</b>	<b>MONOGYNIA.</b>

.....

The original species is a native of Ireland. The present beautiful variety has been in cultivation for a considerable time, constantly retaining its close dwarf character, being rarely above six inches from the ground. It is not known whether it has been found any where in a wild state, or whether it was first raised in a garden from the seeds of the common kind.

It is tolerably hardy, and will thrive either in a pot or border in peat earth. It is propagated freely by cuttings.





*Pleurothallis prolifera.*



No. 1908.

PLEUROTHALLIS PROLIFERA.

Class.	Order.
<i>GYNANDRIA</i>	<i>MONANDRIA.</i>

.....

This is a native of Brazil: according to Mr. Lindley it was received by the Hon. Mr. Herbert, from Boto Fogo, where it was found in shady places, upon steep rocks.

It grows with us very slowly, not having made more than two leaves in four years: the leaves are very concave, and the flowers were produced in November: they lie quite close in the hollow part, in a very curious and unusual manner. But every thing which is created has something in it to excite our admiration of its Glorious Author, that most Beneficent God, who "causeth it to rain on the earth where no man is; on the wilderness where there is no man; to satisfy the desolate and waste grounds, and to cause the bud of the tender herb to spring forth."





*Lactuca scariola*

1847



No. 1909.

LIATRIS GRACILIS.

Class.  
*SYNGENESIA*

Order.  
*ÆQUALIS.*

.....

A native of Georgia: we received it in 1830 from our kind friend Dr. Wray, and it flowered in October 1832. Elliott informs us that it is found growing in dry pine barrens.

It is probably a little tender, therefore it will be prudent to afford it the shelter of a frame in winter. Like the other kinds, it may be increased by dividing the root or by seeds. The soil should be loam and peat.







No. 1910.

**SALVIA FULGENS.**

Class.	Order.
<i>DIANDRIA</i>	<i>MONOGYNIA.</i>

.....

This is a native of Mexico, where it is found on mountains : it was introduced in 1829, and with us it flowers in the month of September and sometimes October.

It is necessary to preserve it in winter in a greenhouse, but being a showy free-growing plant, it is very suitable for planting out in summer ; it will then display its beauties to great advantage during the autumnal months, especially should they be free from frost, which of course would destroy it.

It is easily multiplied by cuttings, and will grow in any soil.



April 1833.

Part 192,

*Price Two Shillings and Sixpence,*

TO BE CONTINUED MONTHLY,

OF THE

# Botanical Cabinet,

CONSISTING OF

COLOURED DELINEATIONS

OF

## PLANTS

FROM ALL COUNTRIES:

WITH

A SHORT ACCOUNT OF EACH, DIRECTIONS FOR  
MANAGEMENT, &c.

BY

CONRAD LODDIGES & SONS.

THE PLATES BY

GEORGE COOKE.

*Each Part will contain Ten Plates, and Ten  
Parts will form one Volume.*

London: Published by

JOHN AND ARTHUR ARCH, CORNHILL;

LONGMAN, REES, ORME, BROWN,

AND CO., PATERNOSTER-BOW;

AND C. LODDIGES AND SONS, HACKNEY.

1833.





*Barleria mitis*



No. 1911.

**BARLERIA MITIS.**

Class.	Order.
<i>DIDYNAMIA</i>	<i>ANGIOSPERMIA.</i>

.....

This is a native of India, introduced about 1816. It is a small upright shrub, and flowers freely in spring and autumn.

It must be protected in the stove, is easy of cultivation, and may be propagated by cuttings, which should be planted in loam and peat earth.





C. Loddige del.

*Billbergia zebrina*



No. 1912.

**BILLBERGIA ZEBRINA.\***

<b>Class.</b>	<b>Order.</b>
<b>HEXÁNDRIA</b>	<b>MONOGYNIA.</b>

.....

This elegant plant is a native of Rio de Janeiro, where it is found growing upon trees. With us it flowered in the month of January. It is cultivated without difficulty in the stove, potted in vegetable earth with a portion of sand, and but little water in the winter season. It will sometimes increase by suckers.

\* We have been obliged to reduce the figure to about half its natural size, in order to give a suitable representation of its proportions.





*Humboldtia* ...



No. 1913.

**THUNBERGIA FRAGRANS.**

Class.	Order.
<i>DIDYNAMIA</i>	<i>ANGIOSPERMIA.</i>

.....

A delicate stove climber: it has been long in cultivation, and is well deserving a place, being almost perpetually adorned with its snow white blossoms. It is said to have been introduced in 1796 by the late Lady de Clifford.

It may be increased with facility by cuttings, and should be potted in light loam.





*Maxillaria punctata*



No. 1914.

**MAXILLARIA PUNCTATA.**

Class.	Order.
<i>GYNANDRIA</i>	<i>MONANDRIA.</i>

.....

This is a native of Brazil, and was introduced a few years since. Its flowers come out in November and December: they possess a very delicate and agreeable fragrance.

It requires the usual treatment of these charming plants, with constant stove heat, and will sometimes admit of separation for increase. It should be potted in vegetable earth with moss and pieces of broken pot intermixed.





*Justicia speciosa*



No. 1915.

**JUSTICIA SPECIOSA.**

Class.	Order.
<i>DIANDRIA</i>	<i>MONOGYNIA.</i>

.....

A native of India, growing in forests in the interior of Bengal, where it is very ornamental.

It was first raised in this country by Sir Henry Bunbury, of Barton Hall, Suffolk.

It is necessary to preserve it in the stove. With us it flowers during the winter. It is easily increased by cuttings, and should be potted in rich loam.





*Psidium tiquetrum.*

111



No. 1916.

**PSILOTUM TRIQUETRUM.**

Class.	Order.
<b>CRYPTOGAMIA</b>	<b>LYCOPODINEÆ.</b>

.....

This is a native of the West Indies, also, it is said, of the Mauritius and of New Holland. We once had a plant which lived many years, growing out of a trunk of *Cycas revoluta*.

It requires the stove, and is a singular looking plant. The capsules are produced at various seasons. It may be potted and preserved in the same way as the tropical ferns.





*Oncidium Harrisonii*



No. 1917.

**ONCIDIUM HARRISONIÆ.**

Class.  
**GYNANDRIA**

Order.  
**MONANDRIA.**

.....

A native of Brazil: we received it from our valued friend Mr. F. Warre. When it arrived it had scarcely a remaining spark of life, but by little and little it has recovered, and produced its graceful flowers in October and November 1832. It requires the heat of the stove, and appears to flourish in a pot, with peat and vegetable earth, intermixed with fragments of pot. It was named by Mr. Lindley, after Mrs. Harrison of Liverpool.

This elegant plant is another choice example of the power and goodness of its Divine Author, of whom indeed "we may speak much and yet come short; wherefore in sum He is All. How shall we be able to magnify Him? For He is great above all His works. The Lord is terrible and very great, and marvellous is His power. When you glorify the Lord, exalt Him as much as you can; for even yet will He far



exceed ; and when you exalt Him, put forth all your strength, and be not weary, for you can never go far enough. Who hath seen Him, that he might tell us? and who can magnify Him as He is? There are yet hid greater things than these be, for we have seen but a few of His works : for the Lord hath made all things.”





*Erica periciliata.*



No. 1918.

**ERICA PENICILLATA.**

Class.	Order.
<i>OCTANDRIA</i>	<i>MONOGYNIA.</i>

.....

This pleasing species is a native of the Cape of Good Hope. It has an affinity to the *Plukenetiana*, but sufficiently distinct. We raised it originally in 1792, and after a time lost it for many years. In 1829 we obtained a few seeds of it again from its native country, from the produce of which our present drawing was made in December 1832. It flowers at various seasons, and ought to have the usual airy greenhouse treatment. It is exceedingly difficult to strike by cuttings, consequently, if it does not ripen its seeds, which is very unlikely, will probably be soon again lost.





*Fraxinus excelsior*



No. 1919.

**LUCULIA GRATISSIMA.**

*Class.*

*Order.*

**PENTANDRIA**

**MONOGYNIA.**

.....

A native of Napal and Silhet, originally classed with Cinchona. According to Dr. Carey, in Flora Indica, it is a small branching tree, growing to the height of sixteen feet, found on the smaller hills in exposed situations, where it flowers nearly all the year. He says it is impossible to conceive any thing more beautiful than this tree, when covered with its numerous rounded panicles of pink, very fragrant, large blossoms.

Our plant flowered in September. It is difficult of cultivation, the stove being too close for it, and the greenhouse too cold. It may be increased by cuttings or layers, and should be potted in loam and peat soil.





*Rhipsalis Mesembryanthemoides*



No. 1920.

**RHIPSALIS MESEMBRYANTHEMOIDES.**

Class.	Order.
<b>ICOSANDRIA</b>	<b>MONOGYNIA.</b>

.....

This curious little plant has been lately introduced from South America. It flowered with us in the month of December, in the stove, which is necessary for its preservation.

It is multiplied without difficulty by cuttings, and should be potted in light loam, with about one-third of decayed mortar.



May 1833.

Part 193,

Price Two Shillings and Sixpence,

TO BE CONTINUED MONTHLY.

OF THE

**Botanical Cabinet,**

CONSISTING OF

COLOURED DELINEATIONS

OF

**PLANTS**

FROM ALL COUNTRIES:

WITH

A SHORT ACCOUNT OF EACH, DIRECTIONS FOR

MANAGEMENT, &c.

BY

**CONRAD LODDIGES & SONS.**

THE PLATES BY

**GEORGE COOKE.**

*Each Part will contain Ten Plates, and Ten  
Parts will form one Volume.*

London: Published by

JOHN AND ARTHUR ARCH, CORNHILL;

LONGMAN, REES, ORME, BROWN,

AND CO., PATERNOSTER-ROW;

AND C. LODDIGES AND SONS, HACKNEY.

1833.





*Justicia callitricha*



No. 1921.

**JUSTICIA CALLITRICHIA.**

Class.	Order.
<i>DIANDRIA</i>	<i>MONOGYNIA.</i>

.....

This singular species is believed to be from Brazil: it was received at the Liverpool Garden in 1826, from Mr. Otto, of Berlin.

It flowers during the winter season, and constantly requires the warmth of the stove, in which it is very ornamental during the dull period of the year.

It may be increased by cuttings, and should be potted in light loam.





1841. 10.

*Mimulus lewisii* Pursh



No. 1922.

**JASMINUM SAMBAC** *pleno*.

Class.	Order.
<b>DIANDRIA</b>	<b>MONOGYNIA.</b>

.....

This is a native of India, where it is much cultivated for the delicious fragrance of its pure white flowers, which are produced in succession for a great part of the year. They are used by the Hindoos in their ceremonies, and are universally esteemed.

With us it requires the stove: it flowers freely at various seasons, and is well deserving a place.

It may be increased by cuttings, and should be potted in rich loam.





*V. Lindley del.*

*Zygopetalum stenochilum*

*J. G. L.*



No. 1923.

**ZYGOPETALUM STENOCHILUM.**

Class.  
***GYNANDRIA***

Order.  
***DIANDRIA.***

... ..

We received this fine plant in 1828, from Mr. Warre, who discovered it and sent it home from Brazil. It flowered with us in the month of October. The blossoms are of long duration, like the others of this beautiful genus, of which there appear to be many kinds, differing slightly from each other, but all very beautiful.

It thrives in the stove potted in vegetable earth, intermixed with broken pieces of pots, and will occasionally increase by offsets.





*Maranta bicolor* Kunth



No. 1924.

**MARANTA BICOLOR** *minor*.

Class.	Order.
<b>MONANDRIA</b>	<b>MONOGYNIA.</b>

.....

This is a native of South America: we received it a few years since from our good friend, Mr. Shepherd, of Liverpool. It is a neat little stove plant, not growing much more than six inches high. The leaves are very richly and beautifully tinted, and the simple flowers are produced in February and March. It should be potted in rich loam, and will increase freely by separating the roots.





*Erigeron annuus*



No. 1925.

**EPACRIS CAMPANULATA.**

<i>Class.</i>	<i>Order.</i>
<b><i>PENTANDRIA</i></b>	<b><i>MONOGYNIA.</i></b>

.....

We raised this elegant plant from New Holland seeds in 1830. It flowered in February and March 1833, and we think does not yield in beauty to any of this fine family.

It will increase by cuttings, and should be kept in the greenhouse, and potted in sandy peat earth.



*Primula sinensis* 1/1



No. 1926.

**PRIMULA SINENSIS** *alba*.

**Class.**

**Order.**

**PENTANDRIA**

**MONOGYNIA.**

.....

This is a variety raised from seeds of the *Primula Sinensis*, differing from the original kind in the colour of the flowers, and the leaves being more deeply divided. It is very showy, and producing its flowers in great abundance in January and February, is very ornamental in the greenhouse.

It is readily raised from seeds, and should be potted in rich light soil.



*Daphne odora.*



No. 1927.

**DAPHNE ODORA.**

Class.	Order.
<i>OCTANDRIA</i>	<i>MONOGYNIA.</i>

.....

This is a native of China: it has been long cultivated in England, and well deserves a place in every greenhouse. Its foliage is glossy and rich, and the flowers are produced very early in the year: their odour is peculiarly grateful, which adds much to its other attractions.

It may be increased by cuttings, and should be potted in loam and peat earth.



Acacia dealbata



No. 1928.

**ACACIA DEALBATA.**

Class.	Order.
<i>POLYGAMIA</i>	<i>MONŒCIA.</i>

.....

A native of New Holland: it has been cultivated for several years, and flowers in February and March. It is fragrant, and the whole plant, with its delicate foliage and splendid flowers, is exceedingly ornamental. It grows freely, and is well adapted for a conservatory.

It is increased by seeds, which are often brought from its native country, where its branches and bark are used in tanning leather.

The soil should be loam and peat.



*Wrightia pubescens*



No. 1929.

**WRIGHTIA PUBESCENS.**

**Class.**

**Order.**

***PENTANDRIA***

***MONOGYNIA.***

.....

This is a native of the India islands, and also of the Tropical parts of New Holland. It is a small shrub, with loose branches, flowering with us in February and March.

It requires the protection of the stove, and may be increased by cuttings or layers, which should be potted in loam and peat earth.



*Acrotriche ovalifolia*



No. 1930.

**ACROTRICHE OVALIFOLIA.**

Class.	Order.
<i>PENTANDRIA</i>	<i>MONOGYNIA.</i>

.....

This was discovered by Dr. Brown on the south coast of New Holland, and has been lately introduced. It is a very low shrub, not exceeding a few inches in height, and flowers in February and March. It may be increased by cuttings, and should be potted in sandy peat soil, and preserved in the greenhouse.

At first sight the flowers of this curious plant appear so minute as to excite little interest, but when we avail ourselves of the help of a magnifying glass, they are found to be exceedingly beautiful in form and the arrangement of their parts. Thus many of the wonderful works of our Glorious Creator do indeed remain hidden, from the weakness of our vision, but are not the less replete with proofs of His infinite skill and transcendant power, which are as marvelous in these minute objects as in those of greater magnitude, and stupendous in all!!

June 1833.

Part 194,

Price Two Shillings and Sixpence,  
TO BE CONTINUED MONTHLY,

OF THE  
**Botanical Cabinet,**

CONSISTING OF  
COLOURED DELINEATIONS

OF  
PLANTS

FROM ALL COUNTRIES:

WITH  
A SHORT ACCOUNT OF EACH, DIRECTIONS FOR  
MANAGEMENT, &c.

BY  
CONRAD LODDIGES & SONS.

THE PLATES BY  
GEORGE COOKE.

*Each Part will contain Ten Plates, and Ten  
Parts will form one Volume.*

London: Published by  
JOHN AND ARTHUR ARCH, CORNHILL;  
LONGMAN, REES, ORME, BROWN,  
AND CO., PATERNOSTER-ROW;  
AND C. LODDIGES AND SONS, HACKNEY.

1833.





*Epacris campamulata alba.*

No. 1931.

**EPACRIS CAMPANULATA** *alba*.

—	Class.	Order.
	<b>PENTANDRIA</b>	<b>MONOGYNIA.</b>

.....

This is a native of New South Wales. We raised it from seeds in 1830, and it flowered in March 1833. It is a very lively and beautiful plant, growing to the height of from one to two feet, and requiring the protection of an airy greenhouse. It should be potted in sandy peat soil, and will propagate by cuttings.





*Melastoma Malabathrica*

No. 1932.

**MELASTOMA MALABATHRICA.**

Class.	Order.
<i>DECANDRIA</i>	<i>MONOGYNIA.</i>

.....

A native of India, introduced about 1793. It is a moderately-sized branching shrub, producing its flowers at various seasons: our drawing was made in March.

It requires the heat of the stove, and flourishes best where plenty of air is admitted, especially in summer. It is multiplied without difficulty by cuttings, and should be potted in loam and peat.





*Prockia crucis.*

No. 1953.

**PROCKIA CRUCIS.**

<b>Class.</b>	<b>Order.</b>
<b>POLYANDRIA</b>	<b>POLYGYNIA.</b>

.....

This is a native of the Havannah: it was brought to the Horticultural Society in 1823, by Mr. J. Don. It is a low bushy shrub, and flowers very freely, in long-continued succession: our drawing was made in March. The flowers are neat, but not shewy: they are most agreeably scented, smelling exactly like mignonette.

It requires constant stove heat, and may be increased by cuttings. The soil should be loam and peat.





*Choldota nubicata*



No. 1934.

**PHOLIDOTA IMBRICATA.**

Class.	Order.
<i>GYNANDRIA</i>	<i>MONANDRIA.</i>

....

A native of Napal, where it grows upon trees. It flowered in the Glasgow Garden in 1824, according to Dr. Hooker. It grows very slowly, and continued flowering with us from February to May, or later, in succession. It has been long known by the figure in Hortus Malabarius, vol. xii. t. 24.

It requires the stove, and may be cultivated in a pot in vegetable earth. It will sometimes admit of being increased by dividing the root.





*Dendrobium pulchellum*

No. 1935.

**DENDROBIUM PULCHELLUM.**

Class.	Order.
<i>GYNANDRIA</i>	<i>MONANDRIA.</i>

.....

This superb species is a native of India : we have cultivated it for three or four years. It appears to grow upon trees, and flowered with us for the first time in February and March 1833. The flowers are exceedingly beautiful in form and colour : they remained several days in perfection.

We have found this plant to thrive best in a small pot of moss, kept moist, and suspended from a rafter in the stove : it makes many branches, with numerous little roots pushing out in all directions.





*Hypoxis ramosa.*

No. 1936.

**HYPOXIS RAMOSA.**

Class.

Order.

*HEXANDRIA*

*MONOGYNIA.*

.....

A native of the Cape of Good Hope : we received it some years since. It flowers in June and July. After flowering, the bulb remains dormant, sometimes for one or two years. It appears to be very slow of increase, not producing offsets. The soil should be sandy peat ; and it is necessary to preserve it in the greenhouse.





*Malpighia lucata.*

No. 1937.

**MALPIGHIA FUCATA.**

Class.	Order.
<i>DECANDRIA</i>	<i>TRIGYNIA.</i>

.....

This is supposed to be a native of the West Indies. It grows erect to the height of three or four feet, and flowers at various seasons: our drawing was made in April. The leaves are covered, especially underneath, with numerous stings, nearly half an inch in length, which have their attachment in the middle, and are sharp at both ends: when touched they enter the skin, and produce considerable irritation.

It requires the protection of the stove, and may be propagated by cuttings. The soil should be loam and peat.





*Stypheha tubiflora*

STYPHELIA TUBIFLORA.

<small>Class.</small>	<small>Order.</small>
<i>PENTANDRIA</i>	<i>MONOGYNIA.</i>

.....

This beautiful plant is a native of New South Wales ; and, although it has long been known by descriptions, and to collectors of dried specimens, we never saw it in a living state till we raised it from seeds in 1830. In March 1833, it produced its elegant flowers : the plant was erect, about two feet in height, with many stiff small branches.

It thrives in sandy peat earth, and should be kept in the greenhouse. It will probably increase by cuttings, although it belongs to a difficult family.

Admiration of the works of our glorious Creator, appears to be a principle which is not subject to decay or to satiety ; though it be perpetually indulged by His great goodness, with the enjoyment of new wonders, it is still thirsting and panting after more, and shall be so doubtless till that blessed period, when the Lord shall give us “ to eat of the Tree of Life, which is in the midst of the Paradise of God.”





No. 1939.

**GESNERIA DOUGLASSII.**

Class.	Order.
<i>DIDYNAMIA</i>	<i>ANGIOSPERMIA.</i>

.....

This is a native of Rio Janeiro, where it was discovered in 1824, by the indefatigable traveller whose name it bears; who, with most active zeal and industry, has done, and is doing, so much for botanical science—not merely by accumulating for the dead gardens of dry collectors, but for the living splendour of almost every garden in Europe, through the whole of which the beautiful plants which he first discovered have been disseminated.

Our present subject requires the stove: it flowers in the spring, continuing long in succession. It may be increased by dividing the root, which is tuberous; and should be potted in rich light earth.





Fig. 1.

*Amphispiza bilineata*

No. 1940.

**KENNEDIA LONGIRACEMOSA.**

Class.	Order.
<i>DIADELPHIA</i>	<i>DECANDRIA.</i>

.....

A native of New Holland, first raised by Mr. Rollison. It is a pretty climber, and, as Mr. Lindley observes, so similar to *monophylla* as hardly to be deemed a species; yet that inconvenience, in a genus so circumscribed in number, is less than such a very long name as it would have as a variety.

It requires the greenhouse, and flowers in spring; may be increased by cuttings; and will thrive in peat earth and loam.



July 1833.

Part 195,

*Price Two Shillings and Sixpence.*

TO BE CONTINUED MONTHLY,

OF THE

# Botanical Cabinet,

CONSISTING OF

COLOURED DELINEATIONS

OF

## PLANTS

FROM ALL COUNTRIES:

WITH

A SHORT ACCOUNT OF EACH, DIRECTIONS FOR  
MANAGEMENT, &c.

BY

CONRAD LODDIGES & SONS.

THE PLATES BY

GEORGE COOKE.

*Each Part will contain Ten Plates, and Ten  
Parts will form one Volume.*

London: Published by

JOHN AND ARTHUR ARCH, CORNHILL;  
LONGMAN, REES, ORME, BROWN,  
AND CO., PATERNOSTER-ROW;  
AND C. LODDIGES AND SONS, HACKNEY.

1833.



*Berberis buxifolia.*



No. 1941.

**BERBERIS BUXIFOLIA.**

Class.	Order.
<i>HEXANDRIA</i>	<i>MONOGYNIA.</i>

.....

This is a native of the Straits of Magellan, and other southern parts of America, and has been lately introduced by Captain King. It forms a low straggling shrub, and flowered with us in April, protected by a cold frame, but in all probability it will prove to be quite hardy enough for our winters. It may be increased by cuttings or layers, and thrives in light loamy soil.



Cyclamen

1841

1841



No. 1942.

CYCLAMEN REPANDUM.

Class.

Order.

*PENTANDRIA*

*MONOGYNIA.*

.....

This pleasing little plant is a native of Greece, and is said to have been introduced in 1816. It produces its flowers in the month of May: they last a considerable time, and will occasionally bear ripe seeds, by which alone it can be multiplied.

The soil should be rich loam: very old rotten dung is excellent for Cyclamens in general. They should be preserved in a greenhouse, near the glass.



*Erica quadrata*

Adapt. J.



No. 1943.

**ERICA QUADRATA.**

Class.	Order.
<i>OCTANDRIA</i>	<i>MONOGYNIA.</i>

.....

A native of the Cape of Good Hope ; introduced in 1829 by Mr. Lee. It flowered abundantly with us in May. The flowers are very angular in their form, and the opening is perfectly four square, which has suggested its name.

It requires the usual greenhouse protection, and may be increased by cuttings. The soil should be peat.





*Rhododendron campanulatum*



No. 1944.

RHODODENDRON CAMPANULATUM.

Class.	Order.
<i>DECANDRIA</i>	<i>MONOGYNIA.</i>

.....

We raised this beautiful plant in 1825, from seeds. It is a native of Napal, and grows, as we were informed by Dr. Wallich, in elevated situations, so that he had no doubt of its bearing our winters uninjured. This has been confirmed by actual experience during the two last seasons, in which several plants remained perfectly well out of doors with us. The young leaves are at first purplish underneath; they then become milk white, and afterwards, when quite hardened, change to a kind of tan colour. It flowered with us in April, kept in a cold frame.

We have succeeded in propagating it by layers, as well as by cuttings. The soil should be loam and peat.

Few plants are so magnificent in flower as the Rhododendrons; few also are so widely diffused over the globe, though

none have yet been found south of the Equator.

We have great pleasure in being enabled to present our friends with this, which is quite new, feeling also a little partiality for the family, as our late revered parent was the first in this country who had for sale the *Rhododendron ponticum*, (then the only species) which is now so common in every garden, and almost in every window. He also introduced the *Chamæcistus*, *Caucasicum*, and *Chrysanthum*, and lately we have added to the number *Barbatum*, *Camtchaticum*, and *Lepidotum*, with the present, and one other unknown species from *Napal*, none of which have yet flowered with us.





*Cyrtanthus spiralis*

No. 1945.

**CYRTANTHUS SPIRALIS.**


Class.	Order.
<i>HEXANDRIA</i>	<i>MONOGYNIA.</i>

.....

This is a native of South Africa, near Algoa Bay. We received bulbs of it in 1831, and it flowered in March 1833, continuing several weeks.

It requires the protection of a greenhouse, and may be potted in sandy peat earth. We have not yet observed any appearance of increase from it, and probably it will therefore continue to be a scarce plant.





*Andromeda hypnoides.*

No. 1946.

**ANDROMEDA HYPNOIDES.**

Class.	Order.
<i>DECANDRIA</i>	<i>MONOGYNIA.</i>

.....

A native of Canada, as well as Lapland and Siberia. It is a minute but elegant plant, growing among moss, to which it very much assimilates itself. It flowered in April and May. It is somewhat difficult of cultivation, succeeding best potted in black peat earth, covered with a bell-glass, and kept in a shady place. It may be increased by cuttings.





*Oxylobium Pultense*

No. 1947.

**OXYLOBIUM PULTENEÆ.**

<b>Class.</b>	<b>Order.</b>
<b>DECANDRIA</b>	<b>MONOGYNIA.</b>

.....

This is a native of New Holland: we received seeds of it in 1829, and it flowered in May last, being about a foot high. It forms a neat little shrub, with rigid branches. The leaves vary greatly in their insertions, on some of the shoots being in fours, on others in threes, and on others scattered.

It requires the greenhouse, and may be increased by cuttings. The soil should be sandy peat.





*Hemantus pulicaris*

Steph. del.

No. 1948.

**HÆMANTHUS PUNICEUS.**

Class.

**HEXANDRIA**

Order.

**MONOGYNIA.**

.....

This plant has been long in cultivation in this country. It is a native of the Cape of Good Hope, and will live in the greenhouse, but appears to thrive most if preserved in the stove. It flowered in May. The roots will bear dividing occasionally, whereby it may be increased. The soil should be loam and sandy peat.





No. 1949.

**GLADIOLUS WATSONIUS.**

Class.	Order.
<i>TRIANDRIA</i>	<i>MONOGYNIA.</i>

.....

This is a native of the Cape of Good Hope. It grows about a foot in height, and flowers with us in May, planted in a border in front of a greenhouse, without any covering or shelter except the wall, which transmits a little warmth, sufficient to preserve the earth in general from freezing.

It increases by offsets, and the soil should be sandy peat.

Although our climate is so different from that whence this beautiful plant is derived, it nevertheless grows and flowers here in great luxuriance. This facility of plants accommodating themselves to such opposite treatment is a great advantage, admitting of so many more kinds being kept in our gardens. It also is another proof of the great goodness of the Almighty Creator, in adapting His works to such varied situations, and giving them the pro-



perty of thriving in them all ; thus vastly extending these sources of innocent delight, and multiplying occasions for cheerful gratitude, and humble love to our Heavenly Father, for his never-ending beneficence !!





No. 1950.

**CANNA PALLIDA** *var.*

Class.	Order.
<b>MONANDRIA</b>	<b>MONOGYNIA.</b>

.....

Our highly-valued friend, the late Mr. Roscoe, in his splendid work on the Scitamineæ, has figured this plant, and describes it as being from the West Indies. We raised ours from seeds received from Mr. Deppe, at Xalapa. It flowered in the month of May. Like the other Cannas (which it much resembles in habit and growth) it will increase by separating the roots, which ought to be planted in rich loam, and kept in the stove.

Aug. 1833.

Part 196,

*Price Two Shillings and Sixpence,*

TO BE CONTINUED MONTHLY,

OF THE

**Botanical Cabinet,**

CONSISTING OF

COLOURED DELINEATIONS

OF

PLANTS

FROM ALL COUNTRIES:

WITH

A SHORT ACCOUNT OF EACH, DIRECTIONS FOR  
MANAGEMENT, &c.

BY

CONRAD LODDIGES & SONS.

THE PLATES BY

GEORGE COOKE.

*Each Part will contain Ten Plates, and Ten  
Parts will form one Volume.*

London: Published by  
JOHN AND ARTHUR ARCH, CORNHILL;  
LONGMAN, REES, ORME, BROWN,  
AND CO., PATERNOSTER-ROW;  
AND C. LODDIGES AND SONS, HACKNEY.

1833.





No. 1951.

HESPERIS TRISTIS.

Class.	Order.
<i>TETRADYNAMIA</i>	<i>SILICUOSA.</i>

.....

This plant is a native of Germany and Switzerland: it is hardy, but usually considered only biennial. It has been long cultivated. The flowers are very fragrant in the night: in the day they are quite scentless. It is necessary to raise it from seeds, as it requires renewing often. It will grow in any good garden soil, either in a pot or in a border.





*Hyacinthus amurensis*

No. 1952.

**HYACINTHUS AMETHYSTINUS.**

Class.

Order.

**HEXANDRIA**

**MONOGYNIA.**

.....

This pleasing little plant is a native of the South of Europe: it grows about six inches high from the bulb, and flowers in April and May. It should be potted in sandy loam, and may be increased occasionally by offsets.





No. 1953.

**CHORIZEMA SPARTIOIDES.**

Class.	Order.
<i>DECANDRIA</i>	<i>MONOGYNIA.</i>

.....

A native of New Holland : we raised it from seeds in 1832, and it flowered in abundance the following spring. The plants were not more than two or three inches high. The flowers are large, and splendid in colour. We have kept it in the greenhouse, potted in sandy peat earth. Its minuteness has hitherto prevented us from any attempts at propagating it, but it will probably, like others of the same genus, strike by cuttings.





*Elichrysum filiforme.*

215 1/2

No. 1954.

**ELICHRYSUM FILIFORME.**

<i>Class.</i>	<i>Order.</i>
<i>SYNGENESIA</i>	<i>SUPERFLUA.</i>

.....

This is a native of the Cape of Good Hope, introduced about the year 1799. It is a bushy low shrub, and flowers in May. It may be propagated by cuttings freely, and should be potted in sandy peat earth, and preserved in an airy greenhouse. When near flowering care should be taken to prevent any wet from getting into the buds, which invariably spoils them.





No. 1955.

**OROBUS AUREUS.**

Class.	Order.
<i>DIADELPHIA</i>	<i>DECANDRIA.</i>

.....

A native of the northern parts of Europe : it is perennial and of course quite hardy, and produces its flowers in abundance in May. The stems are not more than a foot in height. The root may sometimes be separated for increase, although seeds are by far the more preferable way, and we have no doubt they will occasionally be ripened in this country. It will grow in any good garden soil.





No. 1956.

**CATTLEYA LABIATA.**

Class.	Order.
<i>GYNANDRIA</i>	<i>MONANDRIA.</i>

.....

This superb plant is a native of Brazil, where it was first discovered and sent home by Mr. Swainson. It requires the stove, and usually flowers in the latter part of summer. It will now and then admit of separation, although it multiplies but slowly. The soil should be sandy peat, with moss and a good proportion of small fragments of broken pots. Great care is requisite to keep the wood-lice from it, as without this precaution these destructive vermin will absolutely destroy it.





*Sarracenia flava.*

Wm. H. Bennett

No. 1957.

**SARRACENIA FLAVA.**

Class.	Order.
<i>POLYANDRIA</i>	<i>MONOGYNIA.</i>

.....

A native of Carolina and Florida, growing in swamps. Its leaves, which are often more than a foot in length, form a long attenuated tube: this is frequently found to contain a quantity of dead small flies and gnats. The top or lid of the leaf never closes, yet the insects seem to be effectually imprisoned therein. It must be kept in the greenhouse, in a small pan of water. The soil should be sandy peat. We have sometimes, though rarely, been able to separate the root, and thus make two plants out of one.

This is a most extraordinary plant: the uses of its curiously-formed leaves are not at all understood. But when such poor short-sighted creatures as we are strive to dive into the motives and designs of our Great and Glorious Creator, in His marvellous works, alas! how soon are we lost in astonishment; how sensibly do we im-



mediately feel our own nothingness and incapacity to understand even the simplest of the doings of Almighty Wisdom. We can, indeed, but fall down in silent rapture, and gaze, and wonder, and adore!





No. 1958.

**GNIDIA IMBERBIS.**

Class.	Order.
<i>OCTANDRIA</i>	<i>MONOGYNIA.</i>

.....

This is a native of the Cape of Good Hope: it was introduced in 1792, and flowers most part of the summer. Like most of this family, the flowers are fragrant in the night, but scarcely so at all in the day-time. It requires the greenhouse, and may be readily increased by cuttings. The soil should be sandy peat.





No. 1959.

**SOLANUM CRISPUM.**

Class.	Order.
<i>PENTANDRIA</i>	<i>MONOGYNIA.</i>

.....

A native of Chili, very lately introduced. It has been known for many years, from the figure in the *Flora Peruviana*. With us it flowers in May and June, and is supposed to be hardy enough to bear our winters. It grows fast in almost any soil, and will increase by cuttings without difficulty.





No. 1960.

**ROSA BANKSIÆ** *lutea*.

Class.

Order.

**ICOSANDRIA**

**POLYGYNIA.**

.....

This is a very pretty climbing rose: it was introduced in 1824 by the Horticultural Society, and being very freely increased by cuttings, has already become most extensively diffused throughout the country. It usually flowers earlier than most of the roses, and with the shelter and support of a wall will flourish almost in any soil or situation.

Sept. 1833.

Part 197,

*Price Two Shillings and Sixpence,*

TO BE CONTINUED MONTHLY,

OF THE

# Botanical Cabinet.

CONSISTING OF

COLOURED DELINEATIONS

OF

PLANTS

FROM ALL COUNTRIES:

WITH

A SHORT ACCOUNT OF EACH, DIRECTIONS FOR  
MANAGEMENT, &c.

BY

CONRAD LODDIGES & SONS.

THE PLATES BY

GEORGE COOKE.

*Each Part will contain Ten Plates, and Ten  
Parts will form one Volume.*

London:—Published by  
JOHN AND ARTHUR ARCH, CORNHILL;  
LONGMAN, REES, ORME, BROWN,  
AND CO., PATERNOSTER-ROW;  
AND C. LODDIGES AND SONS, HACEVEY.

1833.





*Erica canthariformis*

No. 1961.

**ERICA CANTHARÆFORMIS.**

Class.	Order.
<i>OCTANDRIA</i>	<i>MONOGYNIA.</i>

.....

A native of the Cape of Good Hope, introduced not long since: it is of upright growth, and flowers in May and June: the flowers in shape resemble a little can. It requires the usual treatment of an airy greenhouse, and should be potted in sandy peat earth. It may be propagated by cuttings.





*Oxalis Brasiliensis.*

1. 2. 3. 4. 5.

No. 1962.

**OXALIS BRASILIENSIS.**

Class.	Order.
<i>DECANDRIA</i>	<i>PENTAGYNIA.</i>

.....

A native of Brazil, roots of which we received in 1829 from Mr. Warre, who collected them himself. It flowered in May 1833, and is a very pleasing plant, as indeed are most of this extensive genus.

We have kept it very well in the greenhouse, potted in light loam and peat : it increases itself freely by offsets.





Solanum ligustrinum.

Del.

No. 1963.

**SOLANUM LIGUSTRINUM.**

Class.	Order.
<i>PENTANDRIA</i>	<i>MONOGYNIA.</i>

.....

This is a native of Chili: we received seeds of it from Mr. Cuming, in 1831, and it flowered in May 1833: it is a free growing shrubby plant, with slender branches, and may be easily preserved in a greenhouse.

It will propagate readily by cuttings, which should be potted in rich loamy soil.





*Ixia curta.*

*Ixia curta*

No. 1964.

**IXIA CURTA.**

Class.	Order.
<b>TRIANDRIA</b>	<b>MONOGYNIA.</b>

.....

This splendid plant is a native of the Cape of Good Hope: it grows to the height of about six inches, and flowers in May and June: we find it to succeed best in a border close to the front wall of a stove, where it has the advantage of the open air, yet is very rarely frozen: the soil should be sandy peat. It increases itself freely by offsets from the bulbs, or by seeds.





F. Lindley del.

*Pimelia sylvestris*

No. 1965.

**PIMELEA SYLVESTRIS.**

Class.  
**DIANDRIA**

Order.  
**MONOGYNIA.**

.....

This is a native of New Holland : it has been lately raised by Mr. Knight : it flowers in May and June, and appears to be a small shrub with few branches.

It requires the protection of an airy greenhouse, and will increase by cuttings or seeds, which appear likely to ripen in this country : the soil should be sandy peat.





Idacei lei

*Pimelia hispida*

No. 1966.

**PIMELEA HISPIDA.**

Class.	Order.
<i>DIANDRIA</i>	<i>MONOGYNIA.</i>

.....

This, like the preceding, was raised in 1830, also by Mr. Knight: its growth and habit appear very similar, and it flowers likewise in May and June. It is from New Holland, and must be preserved in a light airy greenhouse.

It should be potted in sandy peat earth, and may be increased by cuttings or seeds.





il. Loddiges del.

*Cirrhæa viridipurpurea*

No. 1967.

**CIRRHÆA VIRIDIPURPUREA.**

<b>Class.</b>	<b>Order.</b>
<b>GYNANDRIA</b>	<b>MONANDRIA.</b>

....

This curious plant is a native of Brazil, and has been lately introduced. It flowered with us in May: the flowers open all at once, and continue for a considerable time.

It requires the stove, and succeeds very well potted in vegetable earth with moss and fragments of broken pot, increasing occasionally by separating the bulbs.





No. 1968.

**BLETIA HYACINTHINA.**

Class.	Order.
<i>GYNANDRIA</i>	<i>MONANDRIA.</i>

.....

A native of China, introduced in 1802 : its flowers are exceedingly beautiful ; they are produced during the spring months.

It has been usual to keep this plant in the stove, but we find it to grow and thrive in a far superior manner in the greenhouse : it should be potted in vegetable earth with a portion of sand, and increases itself freely by offsets from the bulbs.

The colour of this flower baffles art more than many others, its clearness and brilliancy in the living specimen being absolutely inimitable. But this is only one of the innumerable instances of the utter imperfection of human performances when brought into comparison with the works of our Great and Glorious Creator, the wonders of Whose Wisdom, and Power, and Goodness, are so marvellously displayed through such myriads of beautiful examples, adorning and covering the whole earth !





Hoya Pettsii.

No. 1969.

**HOYA POTTSII.**

Class.	Order.
<i>PENTANDRIA</i>	<i>TRIGYNIA.</i>

.....

This was introduced in 1824, from China, to the garden of the Horticultural Society, by their collector, Mr. Potts, after whom it has received its name. It has been kept constantly in the stove, and flowered in the month of May.

It may be increased without difficulty by cuttings, and should be potted in rich loam.





*Iris Hungarica.*

No. 1970.

IRIS HUNGARICA.

Class.  
*TRIANDRIA*

Order.  
*MONOGYNIA*.

.....

This is a native of Hungary: it has been lately introduced, and is quite hardy of course, with us. It produces its pleasing flowers in the months of April and May, and is well worthy of a place in any garden. There is no difficulty of increasing it by dividing the roots. It will thrive in any good soil, either in a pot or in an open border.



Oct. 1833.

Part 198,

Price Two Shillings and Sixpence,

TO BE CONTINUED MONTHLY,

OF THE

# Botanical Cabinet,

CONSISTING OF

COLOURED DELINEATIONS

OF

## PLANTS

FROM ALL COUNTRIES:

WITH

A SHORT ACCOUNT OF EACH, DIRECTIONS FOR  
MANAGEMENT, &c.

BY

CONRAD LODDIGES & SONS.

THE PLATES BY  
GEORGE COOKE.

*Each Part will contain Ten Plates, and Ten  
Parts will form one Volume.*

London: Published by  
JOHN AND ARTHUR ARCH, Cornhill;  
LONGMAN, REES, ORME, BROWN,  
AND CO., PATERNOSTER-ROW:  
AND C. LODDIGES AND SONS, MARKET.

1833.







No. 1971.

**ANISANTHES CUNONIA.**

Class.	Order.
<b>TRIANDRIA</b>	<b>MONOGYNIA.</b>

.....

This splendid bulbous plant is a native of the Cape of Good Hope : it has been long known and cultivated, but never has become common.

It should be potted in sandy peat earth and kept in a light airy greenhouse, with plenty of water in the growing season, but very little at any other time : it increases itself by offsets from the bulbs.



W. A. P. del.

*Campanula grandiflora.*



No. 1972.

**CAMPANULA GRANDIFLORA.**

Class.	Order.
<i>PENTANDRIA</i>	<i>MONOGYNIA.</i>

.....

This is a native of Siberia and Tartary: it was introduced in 1782, and was formerly common in many collections, but has since been almost wholly lost, as it is very difficult to propagate, and does not bear seeds here.

It flowers in July, and appears to thrive pretty well in light loamy soil, preserved in winter in a frame.





No. 1973.

**ERICA SORDIDA.**

Class.	Order.
<i>OCTANDRIA</i>	<i>MONOGYNIA.</i>

.....

This is a native of the Cape of Good Hope: it was introduced in 1810, and is of free growth, flowering in June and July with us.

It requires the usual airy greenhouse treatment, and is increased with facility by cuttings: the soil should be sandy peat.







No. 1974.

**CAMELLIA JAPONICA** *imbricata*.

Class.	Order.
<b>MONADELPHIA</b>	<b>POLYANDRIA.</b>

.....

A native of China, introduced to the garden of the Horticultural Society in 1824, by their collector, Mr. Parks. It is a free growing kind, with rather lax branches: it flowers early, and is among the finest of the varieties, being very large and double, also in some instances beautifully marked, although this is not always the case. In management it requires the same care as has been recommended for the others, and is also increased by grafting upon the single stock.





No. 1975.

**SOLLYA HETEROPHYLLA.**

Class.	Order.
<b>PENTANDRIA</b>	<b>MONOGYNIA.</b>

.....

This is from the south-west coast of New Holland, and has been lately introduced: it was named by Dr. Lindley in compliment to A. H. Solly, Esq. and is a very pretty climbing greenhouse plant, almost hardy enough to bear the open air.

It flowers in June and July, and may be increased by cuttings or seeds: it should be potted in loam and peat.



*Mimulus roseus*

• *Mimulus roseus*



No. 1976.

**MIMULUS ROSEUS.**

Class.	Order.
<i>DIDYNAMIA</i>	<i>ANGIOSPERMIA.</i>

.....

A native of Northern California, sent in 1831, by Mr. Douglas, to the Horticultural Society: it is perennial, and flowers in July and August, sometimes perfecting its seeds here, by which, as also by cuttings, this elegant plant will increase, and probably soon become universally diffused.

It is well to set the pot in a pan of water, which will make it grow and flower much better: the soil should be light loam.



*Bletia gracilis.*



No. 1977.

**BLETIA GRACILIS.**

Class.	Order.
<i>GYNANDRIA</i>	<i>MONANDRIA.</i>

.....

This is believed to be a native of Mexico: we received it in 1830: it is exceedingly slender in all its proportions; the leaves are of a purplish red colour, and the scape is about a foot in height, producing two or three flowers: ours came out in June.

We have preserved it in the stove, potted in sandy peat and vegetable earth. It will occasionally admit of increase, by separating the bulbs.



*Salpiglossis intertextifolia*



No. 1978.

**SALPIGLOSSIS INTEGRIFOLIA.**

Class.	Order.
<i>DIDYNAMIA</i>	<i>ANGIOSPERMIA.</i>

... ..

This showy plant was raised at the Glasgow Botanical Garden, from seeds received from Buenos Ayres, in the neighbourhood of which it is found wild.

It flowers during great part of the summer, and grows freely either in a pot or in the full ground, increasing by cuttings, and requiring greenhouse protection in winter.



J. Ledebour del<sup>o</sup>

*Sisyrinchium iridifolium*



**SISYRINCHIUM IRIDIFOLIUM.**

Class.	Order.
<b>TRIANDRIA</b>	<b>MONOGYNIA.</b>

.....

This is a native of Chili: we raised it from seeds received from Mr. Cuming, but it had already been introduced.

It produces its flowers in May; they last but a few hours, but a succession of them come out for a considerable time: they sometimes perfect their seeds, and may be increased also by dividing the roots.

It should be preserved in a greenhouse, and potted in light loamy soil.

It has been observed that flowers of short duration are usually splendid in colour; but however we may trace some general rules for this as for other things, still none are universal; exceptions being constantly found, shewing how little we are able to understand about the laws of the beautiful works of God, each of which has some perfection, some excellence in form or colour, or some admirable quality to commend it to our notice, and to excite our gratitude to the Glorious Author of our being, the Fountain of all real unfading Happiness and Joy.





No. 1980.

**LUPINUS LEPIDUS.**

Class.	Order.
<i>DIADELPHIA</i>	<i>DECANDRIA.</i>

.....

A very pretty little perennial species, a native of the North-West of America, introduced by Mr. Douglas to the garden of the Horticultural Society. It flowers during the summer months, and does not often exceed six or nine inches in height.

It produces its seeds in this country, by which it is readily increased, and appears to be quite hardy.

Nov. 1843.

Part 199,

*Price Two Shillings and Sixpence,*

TO BE CONTINUED MONTHLY,

OF THE

**Botanical Cabinet,**

CONSISTING OF

COLOURED DELINEATIONS

OF

**PLANTS**

FROM ALL COUNTRIES:

WITH

A SHORT ACCOUNT OF EACH, DIRECTIONS FOR  
MANAGEMENT, &c.

BY

**CONRAD LODDIGES & SONS.**

THE PLATES BY

**GEORGE COOKE.**

*Each Part will contain Ten Plates, and Ten  
Parts will form one Volume.*

London: Published by  
JOHN AND ARTHUR ARCH, CORNHILL;  
LONGMAN, REES, ORME, BROWN,  
AND CO., PATERNOSTER-ROW;  
AND C. LODDIGES AND SONS, HACKNEY.

1843.





No. 1981.

**FUCHSIA GLOBOSA.**

Class.	Order.
<i>OCTANDRIA</i>	<i>MONOGYNIA.</i>

.....

This has been lately introduced, probably from South America ; but its history is unknown. It produces its beautiful flowers during the whole of the summer, kept in a greenhouse, with plenty of air. It appears to be more robust in its habit than most of the kinds, retaining its leaves better in winter.

It will increase abundantly by cuttings, and grows well in rich garden soil.





*Erica hispida*

*Erica hispida* L.

No. 1982.

**ERICA HISPIDA.**

Class.	Order.
<b>OCTANDRIA</b>	<b>MONOGYNIA.</b>

.....

A native of the Cape of Good Hope: we raised a single plant of it from seeds received thence in 1792, and it flowered within two years afterwards.

It is a slender quick growing kind, making many loose branches, which produce flowers near their tops in June and July.

It requires an airy greenhouse, increases freely by cuttings, and should be potted in sandy peat earth.





*Callistachys retusa*

No. 1983.

**CALLISTACHYS RETUSA.**

Class.	Order.
<i>DECANDRIA</i>	<i>MONOGYNIA.</i>

.....

This is a native of New Holland: it has been lately introduced, and flowered with us in succession from June to August. It appears to grow to three or four feet in height, with a few strong branches, each producing a head of flowers. It requires the greenhouse or conservatory, and is readily propagated by cuttings. The soil should be loam and peat.





*Oncidium pulchellum*.

Hedden 18

No. 1984.

**ONCIDIUM PULCHELLUM.**

<b>Class.</b>	<b>Order.</b>
<b>GYNANDRIA</b>	<b>MONANDRIA.</b>

.....

This delicate plant is a native of Demerara, growing on trees. We have had it several years, but it was not strong enough to bloom till March last. In cultivation the flower stem has been tied up to a stick, but we have no doubt its true mode of growth in the natural situation is hanging down, which is far more elegant.

It continued in flower nearly two months, and has a delicious fragrance. We have preserved it of course constantly in the stove, in a pot with vegetable earth and pieces of broken pot, but have not yet been able to increase it.





*Stanhopea insignis*



No. 1985.

**STANHOPEA INSIGNIS.**

Class.

*GYNANDRIA*

Order.

*MONANDRIA.*

.....

This is a native of South America. Dr. Lindley, who suggested to us the name of *Ceratochilus* to another species, having since discovered that the name had been already applied by Blume to a different plant, has now placed it to the genus *Stanhopea*, as originally constituted by Dr. Hooker, in Curtis's Magazine. It appears to grow naturally upon trees, with the flowers hanging down, as depicted in our reduced figure: the other two flowers are of the natural size. It requires the stove, should be potted in vegetable earth, and flowers in October.

Seldom have we witnessed a more extraordinary subject in the size and form of its flowers, as well as in the curious markings of their different parts; but really both the number and the magnificence of the plants of this class become every day more and more astounding: they actually comprise

a botanical world in themselves. Dr. Lindley, in the first three parts of his elaborate *Genera and Species*, has described upwards of a thousand, and this is to be considered only as the beginning of the work, for new genera and species are constantly pouring in from all quarters.





*Chrysanthemum* [illegible]

No. 1986.

THYSANOTUS JUNCEUS.

Class.	Order.
<i>HEXANDRIA</i>	<i>MONOGYNIA.</i>

.....

A native of New Holland, growing near Port Jackson: it was introduced in 1820, but has been lost for several years, till we received roots and seeds of it again in 1832: some of the former flowered in June and July following: the flowers open only once, remaining for a few hours, but more are produced for a considerable time in succession.

The plant must be kept in an airy greenhouse, potted in sandy peat earth: it does not appear likely to increase, unless seeds will ripen, which has not hitherto been the case.

The delicacy and splendour of these flowers is surprising; their colour also peculiarly pleasing and attractive; all produced from a little tuft of herbage, which when not in bloom it is difficult to distinguish from common grass. Overflowing with elegance and brilliancy, and exquisite

skill, are all the works of the gracious Creator, in variety constantly new, in beauty always charming, awakening the mind to cheerful praise of our most Beneficent Father, who thus in a thousand and a thousand delightful ways, is soothing the ruggedness of life's toilsome path, and still endearing Himself more and more to the adoring hearts of His children.





*Goodyera repens*

No. 1987.

**GOODYERA REPENS.**

Class.	Order.
<i>GYNANDRIA</i>	<i>MONANDRIA.</i>

.....

This is a native of the northern parts of Europe and America. It is found, according to Sir James Smith, in Scotland, growing in mossy Alpine woods, but is rare.

It may be cultivated in a pot in peat earth, placed under a shady wall, and will sometimes increase by separating the creeping roots. The flowers are produced in June and July.





No. 1988.

**SPIRÆA GRANDIFLORA.**

Class.	Order.
<b>ICOSANDRIA</b>	<b>PENTAGYNIA.</b>

.....

This beautiful hardy shrub is a native of Kamtschatka. We raised it from seeds sent to us by Mr. Busch, in 1826. It bears a resemblance to *Spiræa salicifolia*, but is a much finer plant, and the flowers are double the size.

It flourishes in the open ground, in light loamy soil, and flowers in July. May be increased by layers or cuttings.



*Trorolix pentaphylla* (L.)

No. 1989.

**TROPÆOLUM PENTAPHYLLUM.**

Class.	Order.
<i>OCTANDRIA</i>	<i>MONOGYNIA.</i>

.....

This has been lately introduced from Buenos Ayres. It has a tuberous root, from whence arises a climbing stem to the height of two feet or more, producing abundance of flowers during the summer months. It requires the greenhouse protection, may be increased by cuttings, and should be potted in rich loamy soil.





Torenia scabra

Wieg. Bot.

No. 1990.

**TORENIA SCABRA.**

Class.	Order.
<i>DIDYNAMIA</i>	<i>ANGIOSPERMIA.</i>

.....

A native of New Holland, introduced in 1831. It grows rapidly, and forms a good-sized bush, with many succulent branches, each producing plenty of flowers, which continue successively from the month of June till autumn.

It requires a warm greenhouse, increases readily by cuttings, and sometimes bears seeds: the soil should be rich loam.

Dec. 1833.

Part 200,

*Price Two Shillings and Sixpence,*

TO BE CONTINUED MONTHLY,

OF THE

**Botanical Cabinet,**

CONSISTING OF

COLOURED DELINEATIONS

OF

**PLANTS**

FROM ALL COUNTRIES:

WITH

A SHORT ACCOUNT OF EACH, DIRECTIONS FOR  
MANAGEMENT, &c.

BY

**CONRAD LODDIGES & SONS.**

THE PLATES BY

**GEORGE COOKE.**

*Each Part will contain Ten Plates, and Ten  
Parts will form one Volume.*

London:—Published by

JOHN AND ARTHUR ARCH, CORNHILL;

LONGMAN, REES, ORME, BROWN,

AND CO. PATERNOSTER-ROW;

AND C. LODDIGES AND SONS, HACKNEY.

1833.



*NOW COMPLETED,*  
THE  
**BOTANICAL CABINET,**

IN 200 PARTS, FORMING 20 VOLUMES;

CONTAINING

2000 FIGURES OF THE MOST INTERESTING PLANTS  
FROM ALL COUNTRIES,

*Accurately drawn and coloured from Nature.*

ENGRAVED BY G. COOKE.

With a short Account of each, and Directions for its Cultivation,  
Management, Soil, &c.

AND A GENERAL INDEX TO THE WHOLE.

BY C. LODDIGES & SONS.

Quarto, price 5s. each Part.

---

*Subscribers having incomplete Sets are respectfully informed that any  
Volume or Part may be had separately.*

---

LONDON:

PUBLISHED BY J. AND A. ARCH, CORNHILL; LONGMAN, REES,  
ORME, BROWN, AND CO. PATERNOSTER-ROW;  
AND C. LODDIGES AND SONS, HACKNEY.



No. 1991.

**ACONITUM STOERCKIANUM.**

Class.	Order.
<b>POLYANDRIA</b>	<b>TRIGYNIA.</b>

.....

This is a hardy perennial plant, a native of Austria, and has been named after the celebrated physician, Baron von Stoerck. It flowers with us in August, and is a very showy plant. In a pot it grows to a foot and a half in height, but in the full ground will attain to twice or thrice that size. It may be increased by separating the roots, which are knobby, and will grow in any good garden soil.





No. 1992.

**ERICA EWERANA** *pilosa*.

Class.	Order.
<b>OCTANDRIA</b>	<b>MONOGYNIA.</b>

.....

A native of the Cape of Good Hope, introduced about the year 1800. It is a vigorous growing kind, and will attain the height of two or three feet. Its flowers are beautiful; they come out in July and August, and last a considerable time.

It requires the usual airy greenhouse protection, will increase by cuttings, and should be potted in sandy peat earth.



*Erica procumbens*



No. 1993.

**ERICA PROCUMBENS.**

Class.	Order.
<i>OCTANDRIA</i>	<i>MONOGYNIA.</i>

.....

A native of the Cape of Good Hope, introduced a few years since. It produces its pleasing flowers during the summer months. It is necessary to keep it, like the other kinds, in a light airy greenhouse. It will increase by cuttings, and should be potted in sandy peat soil.



No. 1994.

**VILLARSIA CHILIENSIS.**

<i>Class.</i>	<i>Order.</i>
<b>PENTANDRIA</b>	<b>MONOGYNIA.</b>

.....

This is a native of Chili, and has been very lately introduced. It flowers in June, with a stem about a foot in height: the flowers are very pretty; and open a few at a time, in succession, each lasting but a short while.

It will increase by separating the root, and should be potted in rich loam, with a pretty large allowance of water.





No. 1995.

HIBISCUS LILIFLORUS.

Class.	Order.
<i>MONADELPHIA</i>	<i>POLYANDRIA.</i>

.....

This elegant plant, of which there are many varieties, was introduced by Mr. Barclay, a few years since. It grows freely, and flowers during the summer. It is necessary to preserve it in the stove, as it is a native of the Mauritius. It will increase readily by cuttings, and should be potted in loam and peat soil.



*Leucostictis ...*




No. 1996.

LINUM CUMINGII.

Class.	Order.
<i>PENTANDRIA</i>	<i>MONOGYNIA.</i>

.....

This pretty little plant is a native of Chili: it was introduced in 1830, by Mr. Cuming. It is very dwarf in its growth, not much exceeding six inches from the ground; and its brilliant flowers are produced during nearly the whole of the summer. It requires the greenhouse, may be propagated by cuttings or seeds, and should be potted in light loam.



No. 1997.

**LANTANA INVOLUCRATA.**

Class.	Order.
<i>DIDYNAMIA</i>	<i>ANGIOSPERMIA.</i>

.....

A native of the West Indies : it has been long known and cultivated, but never has become very common. It requires the stove, and bears its pretty flowers throughout the spring and summer. There is no difficulty in increasing it from cuttings: the soil should be rich loam.





*Leptospermum umbellatum*

No. 1998.

**LEPTOSPERMUM AMBIGUUM.**

Class.	Order.
<i>ICOSANDRIA</i>	<i>MONOGYNIA.</i>

.....

This is a native of New South Wales, and was among the early introductions from that productive country. It is something taller in its growth than most of the species, and will attain the height of four or five feet, being well furnished with pendulous branches, which, loaded with flowers, are very ornamental. The season of flowering is June and July.

It requires the greenhouse protection, and is increased with facility by cuttings: the soil should be loam and peat.





*Cirrha warreana*



No. 1999.

**CIRRHÆA WARREANA,**

Class.

**GYNANDRIA**

Order.

**MONANDRIA.**

.....

This is a native of Brazil: it was discovered by Mr. Warre, who kindly communicated it to us. It bears a strong resemblance to the other species: they are all highly-interesting and curious plants, well deserving every possible care in cultivation.

It succeeds very well in the stove, planted in moss, with potsherds and a little sandy peat soil. Like the others, it will admit of occasional increase by dividing the bulbs.



No. 2000.

**CYCNOCHES LODDIGESII.**

Class.	Order.
<b>GYNANDRIA</b>	<b>MONANDRIA.</b>

.....

This extraordinary plant is a native of Surinam: it was sent to us in 1830, by our kind friend Mr. Lance; and it flowered in May, and again in the winter of 1832. Dr. Lindley has given it this name, and published it in his excellent work on the Genera and Species of Orchideous Plants.

We have preserved it in the stove, suspended from a rafter, and planted in moss and broken bits of pot; but it has not yet increased.

---

Having been enabled to complete our twentieth volume, and thus to place two thousand plants before the public, our labours are closed; the precarious state of our draughtsman's health not permitting him to go on any farther. We cannot take leave of our kind friends without expressing our sincere gratitude for the



encouragement with which they have favoured us in our humble attempts to illustrate a small portion of a most attractive branch of Natural History. We have been richly rewarded for all our exertions by the delight which we have enjoyed, while studying the charming productions of our Almighty Father, so inconceivable in variety, so inimitable in beauty. We know not how to conclude better than in the words of an eloquent writer: "Whatever the department of nature may be which we explore, in quest of evidence for the perfections of its Author, there is no inquirer, though even of the most transcendant powers, who shall ever attain the satisfaction of having traversed the whole length and breadth of the land. He will but have entered and proceeded a certain way, within the margin of a territory, whose riches are inexhaustible."

# INDEX TO VOL. XX.

	No.
<i>Acacia dealbata</i> - - -	<i>Whitish-leaved Acacia</i> 1928
<i>Aconitum Stoereckianum</i> -	<i>Stoerck's Wolfsbane</i> - 1991
<i>Acrotriche ovalifolia</i> - -	<i>Oval-leaved Acrotriche</i> 1930
<i>Andromeda Hypnoides</i> - -	<i>Moss Andromeda</i> - - 1946
<i>Anisanthus Cunonia</i> - -	<i>Cunon's Anisanth</i> - - 1971
<i>Barleria mitis</i> - - -	<i>Yellow Barleria</i> - - - 1911
<i>Berberis buxifolia</i> - - -	<i>Box-leaved Berberry</i> - 1941
<i>Billbergia zebrina</i> . - -	<i>Striped Billbergia</i> - - 1912
<i>Blechnum gracile</i> - - -	<i>Slender Blechnum</i> - - 1905
<i>Bletia gracilis</i> - - -	<i>Slender Bletia</i> - - - 1977
. . . . <i>Hyacinthina</i> - -	<i>Hyacinth-like Bletia</i> - 1963
<i>Callistachys retusa</i> - - -	<i>Blunt-leaved Callista-</i> <i>chys</i> - - - - - 1983
<i>Camellia japonica imbricata</i> - - -	<i>Imbricate Camellia</i> - 1974
<i>Campanula grandiflora</i> -	<i>Great Bell-flower</i> - - 1972
<i>Canna pallida, var.</i> - -	<i>Pale Canna</i> - - - - 1950
<i>Cattleya labiata</i> - - -	<i>Dark-lipped Cattleya</i> - 1956
<i>Chorizema spartioides</i> -	<i>Broom-like Chorizema</i> 1953
<i>Cirrhæa viridiparpurea</i> -	<i>Green purple Cirrhæa</i> 1967
. . . . <i>Warreana</i> - -	<i>Mr. Warre's do.</i> - - - 1999
<i>Clivia nobilis</i> - - -	<i>Noble Clivia</i> - - - - 1906
<i>Cyclamen repandum</i> - -	<i>Repand Cyclamen</i> - - 1942
<i>Cycnoches Loddigesii</i> -	<i>Loddiges's Cycnoches</i> - 2000
<i>Cyrtanthus spiralis</i> - -	<i>Spiral Cyrtanthus</i> - - 1945
<i>Daphne odora</i> - - -	<i>Sweet-scented Daphne</i> 1927
<i>Dendrobium pulchellum</i> -	<i>Beautiful Dendrobium</i> 1935
<i>Elichrysum filiforme</i> - -	<i>Filiform Elichrysum</i> - 1954
<i>Epacris campanulata</i> - -	<i>Bell-flowered Epacris</i> - 1925
. . . . . <i>alba</i>	<i>White do.</i> - - - - - 1931
<i>Erica cantharæformis</i> -	<i>Can-flowered Heath</i> - 1961
. . . <i>Ewerana pilosa</i> - -	<i>Pilose Ewer's do.</i> - - 1992
. . . <i>hispida</i> - - -	<i>Hispid do.</i> - - - - - 1982
. . . <i>penicillata</i> - - -	<i>Pencil-flowered do.</i> - 1918
. . . <i>procumbens</i> - - -	<i>Procumbent do.</i> - - - 1993
. . . <i>quadrata</i> . - -	<i>Square-flowered do.</i> - 1943
. . . <i>sordida</i> - - -	<i>Dull red do.</i> - - - - 1973
<i>Euphorbia punicea</i> - -	<i>Scarlet Euphorbia</i> - - 1901
<i>Fuchsia globosa</i> - - -	<i>Globe-flowered Fuchsia</i> 1981
<i>Gesneria Douglassii</i> . -	<i>Douglas's Gesneria</i> - 1989



INDEX TO VOL. XX.

	No.
Gladiolus Watsonius - - -	<i>Watson's Cornflag</i> - 1949
Gnidia imberbis - - -	<i>Beardless Gnidia</i> - 1958
Goodyera repens - - -	<i>Creeping Goodyera</i> - 1987
Hæmanthus puniceus - - -	<i>Red Blood-flower</i> - 1948
Hedychium stenopetalum	<i>Narrow-petalled Hedy-</i>
	<i>chium</i> - - - - 1902
Hesperis tristis - - -	<i>Night smelling Hesperis</i> - - - - 1951
Hibiscus Liliiflorus - - -	<i>Lily-flowered Hibiscus</i> 1995
Hoya Pottsii - - - - -	<i>Potts's Hoya</i> - - - 1969
Hyacinthus Amethystinus	<i>Amethystine Hyacinth</i> 1952
Hypoxis ramosa - - -	<i>Branching Hypoxis</i> - 1936
Iris Hungarica - - - - -	<i>Hungarian Iris</i> - - 1970
Ixia curta - - - - -	<i>Short Ixia</i> - - - - 1964
Jasminum sambac pleno -	<i>Double Arabian Jas-</i>
	<i>mine</i> - - - - 1922
Justicia callitricha - - -	<i>Beautiful haired Justi-</i>
	<i>cia</i> - - - - 1921
. . . . . speciosa - - -	<i>Shevy do.</i> - - - - 1915
Kennedia longiracemosa -	<i>Long racemed Kenne-</i>
	<i>dia</i> - - - - 1940
Lantana involucrata - - -	<i>Involucrate Lantana</i> - 1997
Leptospermum ambiguum	<i>Ambiguous Leptosper-</i>
	<i>mum</i> - - - - 1998
Liatris gracilis - - - - -	<i>Slender Liatris</i> - - 1909
Linum Cumingii - - - - -	<i>Cuming's Flax</i> - - 1996
Luculia gratissima - - -	<i>Grateful Luculia</i> - 1919
Lupinus lepidus - - - - -	<i>Pretty Lupine</i> - - - 1980
Malpighia fucata - - - - -	<i>Painted Barbadoes</i>
	<i>Cherry</i> - - - - 1937
Maranta bicolor minor -	<i>Lesser two-coloured</i>
	<i>Maranta</i> - - - - 1924
Maxillaria ochroleuca -	<i>Yellowish Maxillaria</i> - 1904
. . . . . punctata - - -	<i>Spotted do.</i> - - - - 1914
Melastoma Malabathrica -	<i>Malabar Melastoma</i> - 1932
Menziesia polifolia nana -	<i>Dwarf Menziesia</i> - - 1907
Mimulus roseus - - - - -	<i>Rose-coloured Monkey-</i>
	<i>flower</i> - - - - 1976
Oncidium Harrisoniæ -	<i>Mrs. Harrison's Onci-</i>
	<i>dium</i> - - - - 1917
. . . . . pulchellum -	<i>Neat Oncidium</i> - - - 1984



INDEX TO VOL. XX.

	No.
<i>Orobus aureus</i> - - - -	<i>Gold-green Vetch</i> - - 1955
<i>Oxalis Braziliensis</i> - -	<i>Brazilian Wood Sor-</i> <i>rel</i> - - - - - 1962
<i>Oxylobium Pulteneæ</i> - -	<i>Pultenea-like Oxylo-</i> <i>bium</i> - - - - - 1947
<i>Pholidota imbricata</i> - -	<i>Imbricate Pholidota</i> - 1934
<i>Pimelea hispida</i> - - - -	<i>Hairy-flowered Pime-</i> <i>lea</i> - - - - - 1966
. . . . . <i>sylvestris</i> - - - -	<i>Wood do.</i> - - - - - 1965
<i>Pleurothallis prolifera</i> -	<i>Proliferous Pleuro-</i> <i>thallis</i> - - - - - 1908
<i>Primula sinensis alba</i> -	<i>White Chinese Prim-</i> <i>rose</i> - - - - - 1926
<i>Prockia Crucis</i> - - - -	<i>Santa Cruz Prockia</i> - 1933
<i>Psilotum triquetrum</i> - -	<i>Triangular Psilotum</i> - 1916
<i>Rhipsalis Mesembryan-</i> <i>themoides</i> - - - - -	<i>Mesembryanthemum-</i> <i>like Rhipsalis</i> - - 1920
<i>Rhododendron campanu-</i> <i>latum</i> - - - - -	<i>Bell-flowered Rhodo-</i> <i>dendron</i> - - - - - 1944
<i>Rosa Banksiæ lutea</i> - -	<i>Yellow Banksia Rose</i> - 1960
<i>Salpiglossis integrifolia</i> -	<i>Entire-leaved Salpi-</i> <i>glossis</i> - - - - - 1978
<i>Salvia fulgens</i> - - - -	<i>Bright red Sage</i> - - - 1910
<i>Sarracenia flava</i> - - - -	<i>Yellow Sarracenia</i> - - 1957
<i>Sisyrinchium Iridifolium</i>	<i>Iris-leaved Sisyrin-</i> <i>chium</i> - - - - - 1979
<i>Solanum crispum</i> - - - -	<i>Curled Nightshade</i> - 1959
. . . . . <i>ligustrinum</i> - - - -	<i>Privet-leaved do.</i> - - 1963
<i>Sollya heterophylla</i> - -	<i>Various-leaved Sollya</i> 1975
<i>Sparaxis tricolor</i> - - - -	<i>Three-coloured Sparaxis</i> 1908
<i>Spiræa grandiflora</i> - - - -	<i>Large-flowering Spiræa</i> 1988
<i>Stanhopea insignis</i> - - - -	<i>Noble Stanhopea</i> - - 1985
<i>Styphelia tubiflora</i> - - - -	<i>Tube-flowered Styphelia</i> 1988
<i>Thunbergia fragrans</i> - - - -	<i>White Thunbergia</i> - 1913
<i>Thysanotus juncens</i> - - - -	<i>Rush-like Thysanotus</i> 1986
<i>Torenia scabra</i> - - - - -	<i>Rough-leaved Torenia</i> 1990
<i>Tropæolum pentaphyllum</i>	<i>Five-leaved Nasturtium</i> 1989
<i>Villarsia Chilensis</i> - - - -	<i>Chili Villarsia</i> - - - 1994
<i>Wrightia pubescens</i> - - - -	<i>Downy Wrightia</i> - - 1929
<i>Zygopetalum stenochilum</i>	<i>Narrow-petalled Zymo-</i> <i>petalum</i> - - - - - 1923

.....  
**W. WILSON, Printer, 57, Skinner-Street, London.**

# GENERAL INDEX TO THE BOTANICAL CABINET.

	No.		No.
<i>Acacia angustifolia</i> .....	763	<i>Alnus cordifolia</i> .....	1231
<i>armata</i> .....	49	<i>ovata</i> .....	1141
<i>brevifolia</i> .....	1235	<i>Aloe denticulata</i> .....	1350
<i>Brownii</i> .....	1333	<i>expansa</i> .....	1430
<i>calamifolia</i> .....	909	<i>humilis</i> .....	1481
<i>dealbata</i> .....	1928	<i>incurva</i> .....	1504
<i>discolor</i> .....	601	<i>reticulata</i> .....	1354
<i>falcata</i> .....	1115	<i>Alonsoa intermedia</i> .....	1456
<i>graveolens</i> .....	1460	<i>Alstonia venenata</i> .....	1180
<i>hispidula</i> .....	823	<i>Alstrœmeria bicolor</i> .....	1497
<i>hybrida</i> .....	1342	<i>Hookeri</i> .....	1272
<i>impressa</i> .....	1319	<i>ligtu</i> .....	17
<i>Juniperina</i> .....	398	<i>oculata</i> .....	1851
<i>linearis</i> .....	595	<i>ovata</i> .....	1654
<i>linifolia</i> .....	383	<i>pelegrina</i> .....	1295
<i>longifolia</i> .....	678	<i>psittacina</i> .....	1666
<i>lophantha</i> .....	716	<i>pulchella</i> .....	1054
<i>lunata</i> .....	384	<i>salsilla</i> .....	1683
<i>melanoxylon</i> .....	630	<i>tricolor</i> .....	1147
<i>Myrtifolia</i> .....	772	<i>Alyssum Alpestre</i> .....	1565
<i>nigricans</i> .....	313	<i>Alyxia ruscifolia</i> .....	1811
<i>ornithophora</i> .....	1469	<i>Amaryllis advena</i> .....	779
<i>prostrata</i> .....	631	<i>aurea</i> .....	847
<i>pulchella</i> .....	212	<i>calyptrata</i> .....	864
<i>Sophoræ</i> .....	1351	<i>crocata</i> .....	1082
<i>stricta</i> .....	99	<i>psittacina</i> .....	1204
<i>suaveolens</i> .....	730	<i>pulverulenta</i> .....	484
<i>Taxifolia</i> .....	1225	<i>rutila</i> .....	1449
<i>verticillata</i> .....	535	<i>Solandræfolia</i> .....	1200
<i>virgata</i> .....	1246	<i>spectabilis</i> .....	159
<i>undulæfolia</i> .....	1544	<i>Amorpha pubescens</i> .....	689
<i>undulata</i> .....	753	<i>Amsonia salicifolia</i> .....	592
<i>Acer hybridum</i> .....	1221	<i>Amygdalus nana</i> .....	1114
<i>Aceras anthropophora</i> .....	1775	<i>orientalis</i> .....	1137
<i>Achania malvaviscus</i> .....	1155	<i>Sibirica</i> .....	1599
<i>mollis</i> .....	600	<i>Anacampseros rotundifolia</i> .....	591
<i>pilosa</i> .....	829	<i>Anagallis latifolia</i> .....	1896
<i>Achillea rosea</i> .....	1407	<i>Anagyris fœtida</i> .....	740
<i>Aconitum Neubergense</i> .....	1410	<i>Anchusa Italica</i> .....	1383
<i>paniculatum</i> .....	810	<i>Andersonia Sprengelioides</i> .....	263
<i>rostratum</i> .....	203	<i>Andromeda arborea</i> .....	1210
<i>Stoerckianum</i> .....	1991	<i>buxifolia</i> .....	1494
<i>versicolor</i> .....	794	<i>calyculata</i> .....	1464
<i>Acrostichum simplex</i> .....	709	<i>latifolia</i> .....	530
<i>Acrotriche ovalifolia</i> .....	1930	<i>nana</i> .....	862
<i>Adesmia microphylla</i> .....	1692	<i>ventricosa</i> .....	1582
<i>Adiantum reniforme</i> .....	841	<i>Catesbæa</i> .....	1320
<i>Ægiphila Martinicensis</i> .....	132	<i>Chinensis</i> .....	1648
<i>Æsculus flava</i> .....	1280	<i>coriacea rubra</i> .....	672
<i>pavia</i> .....	1257	<i>Hypnoides</i> .....	1946
<i>rubicunda</i> .....	1242	<i>Jamaicensis</i> .....	1873
<i>Agapanthus minor</i> .....	42	<i>Ligustrina</i> .....	1110
<i>Ageratum cœlestinum</i> .....	623	<i>polifolia angus-</i>	
<i>Altonia Capensis</i> .....	682	<i>tifolia</i> .....	1591
<i>Albuca major</i> .....	1191	<i>grandiflora</i> .....	1714
<i>minor</i> .....	800	<i>latifolia</i> .....	546
<i>Aletris farinosa</i> .....	1161	<i>revoluta</i> .....	1725
<i>Allamanda cathartica</i> .....	259	<i>rigida</i> .....	430
<i>Allium pendulinum</i> .....	1087	<i>speciosa</i> .....	551
<i>sub-hirsutum</i> .....	943	<i>Androsace carnea</i> .....	40



GENERAL INDEX.

	No.		No.
Androsace chamæjasme ..	232	Arum Orixense .....	422
villosa .....	188	spirale .....	525
Aneimia collina .....	1675	trilobatum .....	516
lanceolata.....	1416	triphyllum <i>zibrinum</i>	820
Anemone acentipetala .....	1704	Asarum Canadense.....	889
alba .....	322	Asclepias Curassavica .....	349
alpina .....	1617	quadrifolia .....	1258
Halleri.....	940	Salicifolia .....	272
palmata .....	175	tuberosa .....	1297
<i>lutea</i> .....	1660	verticillata .....	1067
pratensis .....	900	Aspalathus Cheropoda .....	316
Ranunculoides ..	556	Asphodelus Creticus .....	915
sylvestris.....	1739	fistulosus .....	1124
Thalictroides .....	964	Tauricus.....	1102
<i>pleno</i> .....	770	Aspidistra lurida .....	1468
Anigozanthos flavida .....	1282	Asplenium ebeneum .....	5
Anisanthus Cunonia .....	1971	fiabellifolium ..	1567
Anomatheca cruenta .....	1357	monanthemum..	1700
Anthericum Alooïdes.....	996	palmatum .....	868
glaucum.....	1580	Zamiæfolium ..	854
semibarbatum ..	330	Aster aculeatus .....	830
Antholiza montana.....	1022	Alpinus <i>ramosus</i> .....	1289
Anthyllis Aspalathi .....	1169	blandus .....	959
Cretica .....	597	pulcherrimus .....	6
erinacea .....	318	Astragalus aristatus .....	1278
montana .....	578	brevifolius .....	1388
Aotus villosa .....	1353	Canadensis .....	372
Apargia Alpina .....	539	depressus.....	680
Aphyllanthes Monspeliensis .....	1503	Leontinus .....	432
Aporum anceps .....	1895	leucophæus .....	111
Aquilegia Alpina.....	657	Monspessulanus	981
Canadensis.....	888	Uralensis .....	58
Arabis albida .....	1459	Astrantia minor .....	93
<i>petræa hastulata</i> ..	1304	Astrapæa Wallichii .....	1445
undulata .....	1710	Astroloma humifusum .....	1554
Aralla hispida .....	1306	Atragene Austriaca .....	250
Arbutus mucronata .....	1848	Capensis .....	866
serratifolia .....	580	Sibirica .....	1353
<i>unedo rubra</i> .....	123	Aubrietia purpurea.....	1706
Ardisia canaliculata .....	1083	Azalea calendulacea .....	1324
colorata .....	465	<i>cupræa</i> .....	1391
crenulata .....	2	<i>coccinea major</i> .....	624
elegans .....	264	Indica .....	275
pyramidata .....	448	<i>alba</i> .....	1253
umbellata .....	531	<i>phænicea</i> .....	1735
Arduinia bispinosa .....	387	<i>purpurea pl.</i>	1461
Areuaria ciliata .....	659	nudiflora <i>rubra</i> .....	51
<i>verna</i> .....	98	<i>tricolor</i> ..	1382
Arethusa bulbosa .....	1170	procumbens .....	762
Aretia Alpina .....	297	Sinensis .....	885
pubescens .....	1273	speciosa .....	1255
vitaliana .....	166	verticillata .....	1632
Argemone grandiflora .....	1546	viscosa <i>pubescens</i> ..	441
Aristea capitata .....	1174	<i>rubrescens</i> ..	1518
Aristolochia sempervirens..	231	Babiana ringens .....	1006
tomentosa .....	641	Bæckia virgata .....	341
Armeria fruticosa .....	1587	Banksia marginata .....	61
Arnica crenata.....	901	oblongifolia .....	241
montana .....	1613	paludosa .....	392
scorpioides .....	913	Baphia nitida .....	367
Arum dracontium .....	1163	Baptisia perfoliata .....	1104
flagelliforme .....	306	tinctoria .....	588
odoram .....	416	versicolor .....	1144
		Barbacenia purpurea .....	1532



GENERAL INDEX.

	No.		No.
Barleria alba	360	Cactus truncatus	1207
mitis	1911	Caladium bicolor	255
purpurea	344	lividum	1500
Bauera humilis	1197	maculatum	608
Rubiæfolia	1313	viviparum	281
Begonia dipetala	1730	Zamiæfolium	1408
picta	571	Calanthe Veratrifolia	958
semperflorens	1439	Calathea longibracteata	1719
spatulata	107	macillenta	1781
suaveolens	69	orbiculata	1879
Ulmifolia	638	violacea	1148
Berberis aquifolium	1718	Calceolaria arachnoidea	1557
Buxifolia	1941	bicolor	1783
glumacea	1701	connata	1807
repens	1847	corymbosa	1025
Bergera Koenigii	1019	plantaginea	1402
Besleria Melittifolia	204	purpurea	1754
pulchella	1028	rugosa	942
Betula rubra	1248	thyrsiflora	1649
Bignonia capreolata	714	Calendula graminifolia	57
gracilis	1705	Calla occulta	12
viridiflora	1026	Callicoma serratifolia	1167
Billbergia bicolor	1819	Callistachys retusa	1983
zebrina	1912	Callistemon lophantha	1302
Blechnum gracile	1905	scabra	1288
lanceola	1592	Calopogon pulchellus	340
Bletia gracilis	1977	Calothamnus clavata	1447
Hyacinthina	1968	quadritida	737
pallida	629	villosa	92
Tankervillie	20	Calotis cuneifolia	825
Blighia sapida	1484	Calycanthus præcox	617
Bocconia frutescens	83	Calytrix glabra	586
Bonatea speciosa	284	Camellia axillaris	675
Borbonia lævigata	247	Euryoides	1493
lanceolata	81	Japonica alba	636
Boronia alata	1833	alba-pleno	269
denticulata	1377	Althæiflora	1794
pinnata	473	Anemoniflora	537
serrulata	997	atrorubens	170
Bossiæa heterophylla	271	carnea	455
lenticularis	1238	Chandlerii	1897
linophylla	174	coccinea	1237
microphylla	656	compacta	1836
rufa	1119	corallina	1586
scolopendrium	1747	crassinervis	1475
Brachysema latifolium	411	fimbriata	1103
undulatum	778	imbricata	1974
Bromelia exudans	801	incarnata	140
Broughtonia sanguinea	793	Knightii	1463
Brucea ferruginea	120	Myrtifolia	354
Brunfelsia Americana	553	Pæonæflora	238
violacea	792	Papaveracea	1128
undulata	388	Pomponia	596
Brunia abrotanoides	355	Pressii	1745
lanuginosa	572	punctata	1825
Brunsvigia falcata	745	Rosa-mundi	1866
Bryophyllum calycinum	877	Rosa-sinensis	1455
Bunchosia polystachya	1360	Rossii	1707
Burchellia Capensis	664	rotundifolia	1112
Cactus gibbosus	1524	rubra	397
grandiflorus	1625	variegata	329
speciosissimus	924	varieg. simplex	1694
stellatus	79	Welbankii	1198



GENERAL INDEX.

	No.		No.
<i>Camellia kissi</i> . . . . .	1815	<i>Cineraria aurantia</i> . . . . .	325
<i>Oleifera</i> . . . . .	1065	<i>Cirrhæa viridipurpurea</i> . . . . .	1967
<i>Sasanqua</i> . . . . .	1275	<i>Warreana</i> . . . . .	1999
<i>pleno-carnea</i> . . . . .	1134	<i>Cistus Alpestris</i> . . . . .	131
<i>Campanula aggregata</i> . . . . .	505	<i>helianthemum fulgens</i> . . . . .	202
<i>barbata</i> . . . . .	788	<i>marifolius</i> . . . . .	670
<i>capillaris</i> . . . . .	1406	<i>Claytonia Virginica</i> . . . . .	643
<i>cervicaria</i> . . . . .	452	<i>Clematis angustifolia</i> . . . . .	918
<i>excisa</i> . . . . .	561	<i>aristata</i> . . . . .	620
<i>grandiflora</i> . . . . .	1972	<i>calycina</i> . . . . .	720
<i>linifolia</i> . . . . .	1267	<i>campaniflora</i> . . . . .	987
<i>pulla</i> . . . . .	554	<i>cirrhusa</i> . . . . .	1806
<i>rhomboidea</i> . . . . .	693	<i>ochroleuca</i> . . . . .	661
<i>Sarmatica</i> . . . . .	581	<i>Clerodendrum squamatum</i> . . . . .	796
<i>Scheuchzeri</i> . . . . .	485	<i>Clethra acuminata</i> . . . . .	1427
<i>thyrsoides</i> . . . . .	1644	<i>Cliffortia arachnoides</i> . . . . .	260
<i>Canarina campanula</i> . . . . .	376	<i>Clivia nobilis</i> . . . . .	1906
<i>Canna coccinea</i> . . . . .	739	<i>Cochlearia Danica</i> . . . . .	1482
<i>excelsa</i> . . . . .	743	<i>grœnlandica</i> . . . . .	45
<i>flaccida</i> . . . . .	562	<i>Cœlogyne fimbriata</i> . . . . .	1425
<i>Iridiflora</i> . . . . .	905	<i>Colebrookia oppositifolia</i> . . . . .	487
<i>latifolia</i> . . . . .	634	<i>Colletia Ephedra</i> . . . . .	1830
<i>limbata</i> . . . . .	449	<i>Columnnea scandens</i> . . . . .	403
<i>lutea</i> . . . . .	646	<i>Combretum purpureum</i> . . . . .	563
<i>pallida var.</i> . . . . .	1950	<i>Commelina tuberosa</i> . . . . .	193
<i>patens</i> . . . . .	1693	<i>undulata</i> . . . . .	1553
<i>pedunculata</i> . . . . .	622	<i>Conanthera bifolia</i> . . . . .	904
<i>Caralluma crenulata</i> . . . . .	1774	<i>Convallaria oppositifolia</i> . . . . .	640
<i>fimbriata</i> . . . . .	1863	<i>stellata</i> . . . . .	1080
<i>Carissa Carandas</i> . . . . .	663	<i>verticillata</i> . . . . .	1108
<i>spinarum</i> . . . . .	162	<i>Convolvulus Jalapa</i> . . . . .	518
<i>Carludovica latifolia</i> . . . . .	1068	<i>Coptis trifolia</i> . . . . .	173
<i>Carmichaelia australis</i> . . . . .	1061	<i>Coreopsis grandiflora</i> . . . . .	1290
<i>Carolinea alba</i> . . . . .	752	<i>lanceolata</i> . . . . .	821
<i>insignis</i> . . . . .	1004	<i>Cornus Canadensis</i> . . . . .	651
<i>Cassia Capensis</i> . . . . .	511	<i>Coronilla Iberica</i> . . . . .	789
<i>pulchella</i> . . . . .	1839	<i>juncea</i> . . . . .	235
<i>Casuarina equisetifolia</i> . . . . .	607	<i>Corræa alba</i> . . . . .	152
<i>Catalpa syringæfolia</i> . . . . .	1285	<i>pulchella</i> . . . . .	1684
<i>Catasetum Claveringii</i> . . . . .	1344	<i>speciosa</i> . . . . .	112
<i>Cattleya Forbesii</i> . . . . .	1152	<i>viridis</i> . . . . .	336
<i>guttata</i> . . . . .	1715	<i>Cortusa Matthioli</i> . . . . .	956
<i>labiata</i> . . . . .	1956	<i>Cotoneaster affinis</i> . . . . .	1522
<i>Ceanothus cœruleus</i> . . . . .	110	<i>frigida</i> . . . . .	1512
<i>Ceratochilus grandiflorus</i> . . . . .	1414	<i>melanocarpa</i> . . . . .	1531
<i>oculatus</i> . . . . .	1764	<i>microphylla</i> . . . . .	1374
<i>Cerbera dichotoma</i> . . . . .	1516	<i>Cotyledon coccinea</i> . . . . .	832
<i>fruticosa</i> . . . . .	326	<i>coruscans</i> . . . . .	1030
<i>laurifolia</i> . . . . .	989	<i>ovata</i> . . . . .	1392
<i>manghas</i> . . . . .	304	<i>Crassula capitata</i> . . . . .	1029
<i>Cereus setosus</i> . . . . .	1887	<i>coccinea</i> . . . . .	486
<i>Ceropegia Africana</i> . . . . .	906	<i>cordata</i> . . . . .	359
<i>Cestrum angustifolium</i> . . . . .	618	<i>Jasminea</i> . . . . .	1040
<i>laurifolium</i> . . . . .	1688	<i>lactea</i> . . . . .	735
<i>Chelone obliqua</i> . . . . .	813	<i>versicolor</i> . . . . .	433
<i>Chionanthus virginica</i> . . . . .	1261	<i>undulata</i> . . . . .	584
<i>Chironia Jasminoides</i> . . . . .	27	<i>Cratægus Arbutifolia</i> . . . . .	201
<i>Chorizema Henchmannii</i> . . . . .	1233	<i>glabra</i> . . . . .	248
<i>nana</i> . . . . .	1002	<i>glandulosa</i> . . . . .	1012
<i>rhombea</i> . . . . .	1619	<i>nigra</i> . . . . .	1021
<i>Spartioides</i> . . . . .	1953	<i>oxyacanthapunicea</i> . . . . .	1363
<i>Chrysanthemum arcticum</i> . . . . .	785	<i>spatulata</i> . . . . .	1261
<i>Chrysocoma comaurea</i> . . . . .	833	<i>spinosissima</i> . . . . .	1100



GENERAL INDEX.

	No.		No.
<i>Crinum Broussonetii</i> .....	688	<i>Daviesia glauca</i> .....	43
<i>cruentum</i> .....	346	<i>linearis</i> .....	1615
<i>defixum</i> .....	362	<i>ulicina</i> .....	44
<i>erubescens</i> .....	31	<i>Delphinium Chinense</i> .....	71
<i>Mauritianum</i> .....	650	<i>tricorne</i> .....	306
<i>scabrum</i> .....	529	<i>Dendrobium Pierardii</i> .....	750
<i>toxicarium</i> .....	669	<i>polystachyon</i> ..	458
<i>Crocus pusillus</i> .....	1454	<i>pulchellum</i> ..	1935
<i>reticulatus</i> .....	1822	<i>squalens</i> .....	1059
<i>Crotalaria cordifolia</i> .....	1158	<i>Dentaria digitata</i> .....	757
<i>floribunda</i> .....	509	<i>Dianella strumosa</i> .....	1136
<i>Laburnifolia</i> .....	1796	<i>Dianthus Alpinus</i> .....	67
<i>Croton lineare</i> .....	481	<i>arboreus</i> .....	459
<i>pennicillatum</i> .....	440	<i>punctatus</i> .....	896
<i>pictum</i> .....	870	<i>Dichorizandra oxypetala</i> ..	1440
<i>Crowea saligna</i> .....	310	<i>picta</i> .....	1667
<i>Cullumia ciliaris</i> .....	302	<i>thyrsiflora</i> ..	1196
<i>Cunila Mariana</i> .....	1205	<i>Dicliptera spinosa</i> .....	1244
<i>Cunonia Capensis</i> .....	826	<i>Didymocarpus Rexii</i> .....	1305
<i>Cuphea multiflora</i> .....	808	<i>Digitalis Canariensis</i> .....	1329
<i>Curculigo Sumatrana</i> .....	443	<i>Dillwynia cinerascens</i> .....	527
<i>Cyanella Capensis</i> .....	732	<i>Ericæfolia</i> .....	1277
<i>Cyclamen Coum</i> .....	108	<i>floribunda</i> .....	305
<i>Hederæfolium</i> ..	992	<i>glaberrima</i> .....	582
<i>Persicum</i> .....	751	<i>Juniperina</i> .....	401
<i>repandum</i> .....	1942	<i>parvifolia</i> .....	559
<i>Cyclopia Genistoides</i> .....	1111	<i>pungens</i> .....	1502
<i>Cyrenoches Loddigesii</i> .....	2000	<i>Diosma acuminata</i> .....	493
<i>Cymbidium Aloifolium</i> .....	967	<i>ambigua</i> .....	461
<i>dependens</i> .....	936	<i>amœna</i> .....	161
<i>lancifolium</i> .....	927	<i>capitata</i> .....	860
<i>Sinense</i> .....	37	<i>ciliata</i> .....	210
<i>Cypella Herbertii</i> .....	1810	<i>crenata</i> .....	404
<i>Cypripedium arietinum</i> .....	1240	<i>Cupressina</i> .....	303
<i>calceolus</i> .....	363	<i>fragrans</i> .....	775
<i>humile</i> .....	130	<i>hirta</i> .....	1122
<i>insigne</i> .....	1321	<i>latifolia</i> .....	290
<i>parviflorum</i> ..	414	<i>Linifolia</i> .....	400
<i>pubescens</i> .....	895	<i>rubra</i> .....	1498
<i>spectabile</i> .....	697	<i>serratifolia</i> .....	373
<i>venustum</i> .....	585	<i>sphærocephala</i> .....	1490
<i>Cyrtanthus angustifolius</i> ..	368	<i>tenuissima</i> .....	1624
<i>obliquus</i> .....	947	<i>Dionæa muscipula</i> .....	48
<i>spiralis</i> .....	1945	<i>Diospyrus Vaccinioides</i> .....	1540
<i>Cyrtopodium Andersonii</i> ..	121	<i>Diplazium plantagineum</i> ..	1688
<i>Cytisus calycinus</i> .....	673	<i>Dodecatheon Meadia alba</i> ..	1489
<i>capitatus</i> .....	497	<i>Dodonæa attenuata</i> .....	1470
<i>falcatus</i> .....	520	<i>Dondia Epipactis</i> .....	1832
<i>nigricans</i> .....	570	<i>Doodia aspera</i> .....	39
<i>proliferus</i> .....	761	<i>Doronicum Caucasianum</i> ..	1465
<i>purpureus</i> .....	892	<i>Dorstenia Arifolia</i> .....	999
<i>Dalibarda fragaroides</i> .....	408	<i>ceratosanthes</i> ..	1216
<i>Daphne Alpina</i> .....	66	<i>drakena</i> .....	677
<i>Altaica</i> .....	399	<i>Houstoni</i> .....	1005
<i>Cneorum</i> .....	1800	<i>infundibuliformis</i> ..	1784
<i>collina</i> .....	1348	<i>Doryanthes excelsa</i> .....	765
<i>Gnidium</i> .....	150	<i>Draba stellata</i> .....	32
<i>Napolitana</i> .....	719	<i>Dracæna ferrea</i> .....	136
<i>odora</i> .....	1927	<i>terminalis</i> .....	1224
<i>Oleoides</i> .....	299	<i>umbraculifera</i> ..	289
<i>Davallia Canariensis</i> .....	142	<i>Dracocephalum Argunense</i> ..	797
<i>Daviesia acicularis</i> .....	1234	<i>denticulatum</i> ..	1400
<i>Genistoides</i> .....	1552	<i>grandiflorum</i> ..	186



GENERAL INDEX.

	No.		No.
<i>Dracophyllum capitatum</i> ..	1846	<i>Erica ardens</i> .....	47
<i>gracile</i> .....	1346	<i>aristata</i> .....	73
<i>Drimia acuminata</i> .....	1041	<i>australis</i> .....	1472
<i>lanceæfolia</i> .....	278	<i>barbata</i> .....	124
<i>undulata</i> .....	1898	<i>Beaumontiana</i> .....	1686
<i>Duranta Plumierii</i> .....	280	<i>Bergiana</i> .....	939
<i>Eccremocarpus scaber</i> .....	1411	<i>bicolor</i> .....	1001
<i>Edwardsia grandiflora</i> .....	1162	<i>biflora</i> .....	683
<i>Eleagnus angustifolia</i> .....	1339	<i>Blæria</i> .....	85
<i>Eleocarpus cyaneus</i> .....	1055	<i>blanda</i> .....	13
<i>Elichrysum fasciculatum</i> ..	331	<i>Blandfordiana</i> .....	115
<i>filiforme</i> .....	1954	<i>Bonplandiana</i> .....	345
<i>proliferum</i> .....	8	<i>Bowieana</i> .....	842
<i>Scsamoides</i> .....	1149	<i>Bruniades</i> .....	1365
<i>spectabile</i> .....	59	<i>bucciniflora</i> .....	1127
<i>Enkianthus quinqueflorus</i> ..	1101	<i>Caffra</i> .....	196
<i>Epacris attenuata</i> .....	38	<i>calostoma</i> .....	1759
<i>campanulata</i> .....	1925	<i>calycina</i> .....	594
<i>alba</i> ..	1931	<i>campanulata</i> .....	184
<i>Diosmæfolia</i> .....	1634	<i>cantharæformis</i> .....	1961
<i>grandiflora</i> .....	21	<i>capitata</i> .....	1519
<i>impressa</i> .....	1691	<i>carinata</i> .....	1071
<i>nivalis</i> .....	1821	<i>carnea</i> .....	1452
<i>obtusifolia</i> .....	292	<i>carniula</i> .....	926
<i>paludosa</i> .....	1226	<i>Celsiana</i> .....	1777
<i>pulchella</i> .....	194	<i>Cerinthoides</i> .....	1679
<i>purpurascens</i> .....	237	<i>cernua</i> .....	822
<i>rubra</i> ..	876	<i>ciliaris</i> ..	1805
<i>rosea</i> .....	863	<i>cinerea atropurpurea</i> ..	1409
<i>variabilis</i> .....	1816	<i>pallida</i> .....	1505
<i>Epidendrum anceps</i> .....	887	<i>Cliffordiana</i> .....	34
<i>ciliare</i> .....	9	<i>coccinea</i> .....	1375
<i>cochleatum</i> ..	22	<i>colorans</i> .....	224
<i>cuspidatum</i> ..	10	<i>concava</i> .....	134
<i>diffusum</i> .....	846	<i>conferta</i> .....	1335
<i>ellipticum</i> .....	1276	<i>congesta</i> .....	1743
<i>elongatum</i> .....	986	<i>conica</i> .....	1179
<i>fragrans</i> .....	1039	<i>Coventryana</i> .....	423
<i>fuscatum</i> .....	472	<i>crinita</i> .....	1432
<i>nocturnum</i> ..	713	<i>cruenta</i> .....	1656
<i>nutans</i> .....	645	<i>cubica</i> .....	972
<i>patens</i> .....	1537	<i>curviflora</i> .....	1663
<i>polybulbon</i> ..	1230	<i>cylindrica</i> .....	1734
<i>rigidum</i> .....	1600	<i>Daphneflora</i> .....	543
<i>verrucosum</i> ..	1084	<i>Daphnoides</i> .....	154
<i>violaceum</i> .....	337	<i>declinata</i> .....	1662
<i>virescens</i> .....	1867	<i>decora</i> .....	1385
<i>umbellatum</i> ..	26	<i>denticulata</i> .....	1000
<i>Epigæa repens</i> .....	160	<i>dichromata</i> .....	1813
<i>Epimedium diphyllum</i> .....	1858	<i>Dickinsoni</i> .....	1793
<i>Epipactis latifolia</i> .....	982	<i>discolor</i> .....	1453
<i>palustris</i> .....	156	<i>Droseroides</i> .....	1645
<i>Eria rosea</i> .....	1817	<i>Echiiflora</i> .....	364
<i>Erica acuminata</i> .....	216	<i>elata</i> .....	1788
<i>acutangula</i> .....	1868	<i>elegans</i> .....	185
<i>aggregata</i> .....	1678	<i>elongata</i> .....	738
<i>Aitoniana</i> .....	144	<i>Empetrifolia</i> .....	1875
<i>albans</i> .....	95	<i>Empetroides</i> .....	1758
<i>alopecuroides</i> .....	874	<i>epistomia</i> .....	1186
<i>ampullacea</i> .....	508	<i>eriocephala</i> .....	1270
<i>andromedæflora</i> .....	521	<i>erosa</i> .....	133
<i>arbuscula</i> .....	843	<i>erubescens</i> .....	1826
<i>Archeriana</i> .....	1466	<i>Ewerana</i> .....	1303

GENERAL INDEX.

	No.		No.
Erica Ewerana <i>pilosa</i> . . . . .	1992	Erica ostrina . . . . .	1218
eximia . . . . .	1105	ovata . . . . .	417
exposita . . . . .	1521	pallida . . . . .	1355
exsudans . . . . .	287	palustris . . . . .	4
exurgens . . . . .	835	paniculata . . . . .	1194
fastigiata . . . . .	207	Parmentierii . . . . .	197
filamentosa . . . . .	395	patens . . . . .	1228
fimbriata . . . . .	1047	Patersoni . . . . .	1727
flava . . . . .	882	pellucida . . . . .	276
flexuosa . . . . .	1495	pendula . . . . .	902
floribunda . . . . .	176	penicillata . . . . .	1918
florida . . . . .	234	petiolata . . . . .	1150
fragrans . . . . .	288	Petiveri . . . . .	1426
fulgida . . . . .	1633	Peziza . . . . .	265
gelida . . . . .	699	physodes . . . . .	223
gemmifera . . . . .	457	pilosa . . . . .	606
gracilis . . . . .	244	pilularis . . . . .	1563
grandiflora . . . . .	498	Pinea <i>purpurea</i> . . . . .	1259
grandinosa . . . . .	627	Plukenetiana . . . . .	1274
hirsuta . . . . .	754	plumosa . . . . .	1702
hirta . . . . .	1116	Pohlmannii . . . . .	1852
hirtiflora . . . . .	1812	præcox . . . . .	1413
hispidula . . . . .	1982	prægnans . . . . .	945
Humeana . . . . .	389	præstans . . . . .	1605
imbricata . . . . .	1243	Primuloides . . . . .	715
incarnata . . . . .	1655	princeps . . . . .	647
infundibuliformis . . . . .	589	procumbens . . . . .	1993
intertexta . . . . .	1034	propendens . . . . .	63
Irbyana . . . . .	816	pubescens . . . . .	167
Julliana . . . . .	799	pulchella . . . . .	307
lactiflora . . . . .	991	pura . . . . .	72
lævis . . . . .	1393	purpurea . . . . .	703
Lambertiana . . . . .	3	pyramidalis . . . . .	319
Lawsoniana . . . . .	488	quadrata . . . . .	1943
Leeana . . . . .	298	ramentacea . . . . .	446
Linnæana . . . . .	102	recurvata . . . . .	1093
<i>superba</i> . . . . .	1778	reflexa . . . . .	1787
longiflora . . . . .	983	regerminans . . . . .	1728
longipedunculata . . . . .	103	resinosa . . . . .	679
lutea . . . . .	64	retorta . . . . .	804
mammosa . . . . .	125	rigida . . . . .	1286
<i>pallida</i> . . . . .	951	rosea . . . . .	782
Massoni . . . . .	1069	rubella . . . . .	658
melanthera . . . . .	867	rubens . . . . .	557
melastoma . . . . .	333	rubida . . . . .	1166
metulæflora . . . . .	1763	rupestris . . . . .	855
mirabilis . . . . .	1804	Russelliana . . . . .	1013
moschata . . . . .	614	sanguinea . . . . .	86
mucosa . . . . .	35	sanguinolenta . . . . .	468
mucosoides . . . . .	1202	Savillii . . . . .	96
multiflora . . . . .	1572	scabriuscula . . . . .	517
mundula . . . . .	114	scariosa . . . . .	477
mutabilis . . . . .	46	Schollii . . . . .	538
nidularia . . . . .	764	Sebana . . . . .	23
nigrita . . . . .	54	<i>lutea</i> . . . . .	266
nitida . . . . .	1131	Serpyllifolia . . . . .	744
obliqua . . . . .	14	serratifolia . . . . .	1703
obtusa . . . . .	1027	Shannoni . . . . .	168
odorata . . . . .	633	Smithiana . . . . .	1614
ollula . . . . .	1646	sordida . . . . .	1973
oppositifolia <i>alba</i> . . . . .	1343	sparsa . . . . .	1467
<i>rubra</i> . . . . .	1060	speciosa . . . . .	575
orbicularis . . . . .	153	spicata . . . . .	1203



GENERAL INDEX.

	No.		No.
Erica spumosa .....	566	Euonymus bullatus .....	1749
stellata .....	893	Euphorbia anacantha .....	220
stellifera .....	1622	canaliculata .....	727
suaveolens .....	24	caput medusæ ..	1315
sulphurea .....	1762	corollata .....	390
tenella .....	375	Cyparissias .....	118
tenuiflora .....	1717	Ipecacuanha ..	1145
tetragona .....	1239	lophogona .....	1477
Thalictriflora .....	1294	meloformis .....	436
Thunbergii .....	277	polygonata .....	1334
transparens .....	177	punicea .....	1901
triceps .....	962	Salicifolia .....	973
triflora .....	1733	splendens .....	1713
triumphans .....	257	Eurya Chinensis .....	1213
trossula .....	668	Euthalis trinervis .....	1106
<i>rubra</i> .....	1742	Fabricia stricta .....	1219
tubiuscula .....	1157	Fernandesia elegans .....	1214
varia .....	1325	Ferraria atrata .....	1356
ventricosa .....	431	Ficus quercifolia .....	1540
verecunda .....	1827	setacea .....	1138
vernalis .....	1608	urophylla .....	1697
vernix .....	1443	Fontanesia Phillyrioides ..	1308
versicolor .....	208	Fothergilla Alnifolia .....	1576
<i>major</i> .....	1316	Gardeni .....	1507
verticillata .....	145	major .....	1520
vestita <i>alba</i> .....	243	Franciscea uniflora .....	1332
<i>blanda</i> .....	1716	Francoa Sonchifolia .....	1864
<i>carnea</i> .....	1696	Frankenia pauciflora .....	1790
<i>coccinea</i> .....	55	Fuchsia coccinea .....	933
<i>purpurea</i> .....	217	excorticata .....	1347
villosiuscula .....	1844	globosa .....	1981
viridescens .....	233	gracilis .....	934
viridiflora .....	917	macrostemon .....	1062
viscaria .....	726	microphylla .....	1545
umbellata .....	1217	multiflora .....	1415
undulata .....	1792	Fumaria cava <i>alba</i> .....	1589
urceolaris .....	1894	nobilis .....	1630
Walkerii .....	256	Galanthus plicatus .....	1823
Erigeron Alpinum .....	590	Galium Græcum .....	1373
glabellum .....	1631	Garcinia mangostana .....	845
Villarsii .....	1390	Gardenia amœna .....	935
Erinus Alpinus .....	969	<i>angustifolia</i> .....	512
Lychnidea .....	957	Rothmannii .....	1053
Eriocaulon decangulare .....	1310	Gastrolobium bilobum .....	70
Eriostemon Buxifolium ..	1831	Gastronema pallidum .....	1808
cuspidatum ..	1247	Gaultheria procumbens .....	82
Eryngium virgatum .....	1636	Shallon ..	1372
Erysimum pumilum .....	899	Gazania uniflora .....	795
Erythrina cristagalli .....	296	Gela lanceolata .....	938
herbaceum .....	851	Genista Canariensis .....	1201
Escallonia floribunda .....	1772	decumbens .....	718
glandulosa .....	1291	Hispanica .....	1738
rubra .....	1659	ovata .....	482
Eschscholtzia Californica ..	1635	scariosa .....	1135
Eucalyptus Persicifolia .....	501	Gentiana Alpina .....	476
cordata .....	328	Bavarica .....	1256
Euchylus obcordatus .....	60	intermedia .....	218
Eugenia Malaccensis .....	555	purpurea .....	583
myrtifolia .....	625	Saponaria .....	815
Zeylanica .....	690	septemfida .....	89
Eulophia gracilis .....	1178	verna .....	62
Guineensis .....	818	Geodorum dilatatum .....	1797
Euonymus Americanus .....	1322	Geranium argenteum .....	945



GENERAL INDEX.

	No.		No.
<i>Gesneria bulbosa</i> .....	1724	<i>Habranthus robustus</i> .....	1761
<i>Douglasii</i> .....	1939	<i>Hæmanthus albiflos</i> .....	602
<i>rutila</i> .....	1801	<i>coccineus</i> .....	240
<i>Geum coccineum</i> .....	1527	<i>multiflorus</i> .....	912
<i>triflorum</i> .....	1609	<i>pubescens</i> .....	702
<i>Gilliesia graminea</i> .....	1755	<i>punicus</i> .....	1948
<i>Gladiolus Natalensis</i> .....	1756	<i>Hakea ferruginea</i> .....	1501
<i>Watsonius</i> .....	1949	<i>microcarpa</i> .....	219
<i>Globba Careyana</i> .....	691	<i>obliqua</i> .....	1682
<i>Marantina</i> .....	100	<i>pugioniformis</i> .....	353
<i>Globularia vulgaris</i> .....	1888	<i>repanda</i> .....	1750
<i>Gloxinia caulescens</i> .....	1566	<i>Halesia diptera</i> .....	1172
<i>hirsuta</i> .....	1296	<i>tetraptera</i> .....	1173
<i>speciosa</i> .....	28	<i>Hallia imbricata</i> .....	381
<i>Glycine Sinensis</i> .....	773	<i>Hamamelis Virginica</i> .....	598
<i>Gnidia imberbis</i> .....	1958	<i>Hamellia chrysantha</i> .....	1098
<i>imbricata</i> .....	890	<i>Hedera capitata</i> .....	946
<i>ochroleuca</i> .....	1184	<i>Hedychium acuminatum</i> ..	1795
<i>oppositifolia</i> .....	16	<i>carneum</i> .....	693
<i>pinifolia</i> .....	7	<i>coccineum</i> .....	705
<i>radiata</i> .....	29	<i>coronarium</i> .....	507
<i>Gomesa recurva</i> .....	660	<i>elatum</i> .....	856
<i>tenuiflora</i> .....	806	<i>ellipticum</i> .....	1881
<i>Gompholobium pedunculare</i> 1639		<i>flavescens</i> .....	723
<i>Gonolobus grandiflorus</i> ....	1538	<i>flavum</i> .....	604
<i>hirsutus</i> .....	365	<i>spicatum</i> .....	653
<i>Goodenia gracilis</i> .....	1032	<i>stenopetalum</i> ..	1902
<i>Goodia Lotifolia</i> .....	696	<i>urophyllum</i> ..	1785
<i>Geodyera discolor</i> .....	143	<i>Hedysarum carneum</i> .....	312
<i>pubescens</i> .....	1	<i>elongatum</i> .....	1401
<i>repens</i> .....	1987	<i>obscurum</i> .....	1434
<i>tesellata</i> .....	952	<i>Heliconia Bihai</i> .....	252
<i>Govenia superba</i> .....	1709	<i>Helicteris Isora</i> .....	119
<i>Grammitis Ceterach</i> .....	15	<i>Verbascifolia</i> ..	504
<i>Gratiola aurea</i> .....	1399	<i>Helonias Asphodeloides</i> ....	394
<i>Grevillea Acanthifolia</i> ....	1153	<i>bracteata</i> .....	1330
<i>Buxifolia</i> .....	1562	<i>bullata</i> .....	961
<i>cinerea</i> .....	857	<i>læta</i> .....	998
<i>Juniperina</i> .....	1008	<i>Hemerocallis lancifolia</i> ....	1658
<i>linearis</i> .....	50	<i>Sieboldtiana</i> ..	1869
<i>alba</i> .....	858	<i>Herbertia pulchella</i> .....	1547
<i>planifolia</i> .....	1737	<i>Hesperis tristis</i> .....	1951
<i>pubescens</i> .....	1229	<i>Heterotaxis crassifolia</i> ....	1657
<i>punicæ</i> .....	1357	<i>Hibbertia dentata</i> .....	347
<i>Rosmarinifolia</i> ..	1479	<i>Grossularisefolia</i> -	879
<i>sericea</i> .....	880	<i>pedunculata</i> ....	1139
<i>sulphurea</i> .....	1723	<i>Hibiscus Liliiflorus</i> .....	1995
<i>Grewia occidentalis</i> .....	1526	<i>Lindleyi</i> .....	1849
<i>Guatteria rufa</i> .....	612	<i>pedunculatus</i> ....	836
<i>Guzmania tricolor</i> .....	462	<i>Rosa-Malabarica</i> -	769
<i>Gypsophila glomerata</i> ....	1384	<i>Rosa-Sinensis flava</i> 513	
<i>Habenaria albida</i> .....	1121	<i>lutea</i> ....	932
<i>bifolia</i> .....	1653	<i>rubra</i> ..	995
<i>blephariglottis</i> ..	925	<i>variegata</i> 963	
<i>ciliaris</i> .....	1326	<i>splendens</i> .....	1835
<i>cristata</i> .....	1661	<i>Homalium racemosum</i> ....	261
<i>fimbriata</i> .....	552	<i>Hovea Celsi</i> .....	1488
<i>lacera</i> .....	229	<i>elliptica</i> .....	1450
<i>orbiculata</i> .....	1623	<i>latifolia</i> .....	30
<i>tridentata</i> .....	1637	<i>linearis</i> .....	1222
<i>Habranthus Andersonii</i> ....	1677	<i>longifolia</i> .....	994
<i>pallidus</i> .....	1760	<i>purpurea</i> .....	1457
<i>pumilus</i> .....	1771	<i>Houstonia corulea</i> .....	950



GENERAL INDEX.

	No.		No.
Houstonia purpurea	1621	Kæmpferia galanga	980
Hoya pallida	1309	pandurata	587
Pottsii	1969	Kalmia angustifolia <i>rubra</i>	502
Hudsonia Ericoides	192	glauca	1508
Hunnemania Fumariæfolia	1779	hirsuta	1058
Hutchinsia Alpina	766	Kennedia coccinea	1126
Hyacinthus Amethystinus	1952	longiracemosa	1940
Hyoscyamus Orientale	1345	monophylla	758
Hypericum Canariensis	953	ovata	1007
Hypoxis alba	1074	prostrata	1046
erecta	710	rubicunda	954
ramosa	1936	Kleinia suffruticosa	1561
scabra	970	Knowltonia rigida	850
stellata	802	Lachenalia bicolor	1129
Iberis Tenoreana	1721	bifolia	920
Ilex perado	549	fragrans	1140
Illicium Floridanum	209	luteola	734
Indigofera Australis	149	mutabilis	1076
denudata	500	pendula	267
Iris bicolor	1886	quadricolor	746
crassifolia	1861	tricolor	767
cristata	1366	Lachnæa purpurea	273
Hungarica	1970	Lagerstrœmia Indica <i>rosea</i>	1765
Moræoides	1423	Lambertia formosa	80
Nertchinskia	1843	Lantana involucrata	1997
pumila	1574	Lavandulæfolia	1573
reticulata	1829	mista	68
Ruthenica	1650	scabrida	1171
Taurica	1506	Lasiopetalum dumosum	1564
verna	1855	parviflorum	413
Isochilus linearis	1341	purpureum	361
Isopogon Anemonefolius	1337	Quercifolium	619
Isotoma axillaris	1528	Solanaceum	279
Ixia curta	1964	Laurus Cinnamomum	91
patens	1142	Lechenaultia formosa	1066
viridiflora	1548	multiflora	1579
Ixora arborea	609	Ledum Buxifolium	52
coccinea	396	Canadense	1049
crocata	966	latifolium	534
cuneifolia	1215	palustre	500
grandiflora	314	Leontice Thalictroides	1473
incarnata	1048	Leptospermum ambiguum	1998
rosea	729	grandiflorum	514
Jacksonia scoparia	427	grandifolium	701
Jasione perennis	923	lanigerum	1192
Jasminum paniculatum	469	triloculare	791
revolutum	966	Leucocoryne odorata	1768
sambac <i>pleno</i>	1922	Leucojum autumnale	812
Jatropha Gossipifolia	117	pulchellum	1478
panduræfolia	1604	Leucopogon interruptus	1451
urens	478	Juniperinus	447
Jeffersonia diphylla	1036	polystachyus	1436
Justicia asperula	1681	Liatris gracilis	1909
bicolor	166	pilosa	356
callitricha	1921	pumila	147
coccinea	903	scariosa	944
gendarussa	1505	sphaeroldea	1417
maculata	626	Lightfootia tenella	1038
nervosa	146	Lilium autumnale	335
Orchioides	827	Buschianum	1628
retusa	724	Catesbæa	807
speciosa	1915	croceum	784
vanusta	1791	Japonicum	438



GENERAL INDEX.

	No.		No.
<i>Lilium longiflorum</i> .....	985	<i>Malva umbellata</i> .....	222
<i>Philadelphicum</i> .....	976	<i>Mammillaria discolor</i> .....	1671
<i>pumilum</i> .....	358	<i>Mantisia saltatoria</i> .....	137
<i>Limodorum maculatum</i> .....	496	<i>Maranta bicolor</i> .....	921
<i>Limonia trifoliata</i> .....	18	<i>minor</i> .....	1924
<i>Linaria Alpina</i> .....	415	<i>zebrina</i> .....	494
<i>Genistifolia</i> .....	380	<i>Marica cœrulea</i> .....	407
<i>triornithophora</i> ..	1010	<i>humilis</i> .....	1081
<i>Linnæa borealis</i> .....	183	<i>Martinicensis</i> .....	1362
<i>Linum angustifolium</i> .....	1543	<i>Northiana</i> .....	873
<i>campanulatum</i> .....	1254	<i>Sabini</i> ..	1164
<i>Cumingii</i> .....	1996	<i>semi-aperta</i> .....	685
<i>montanum</i> .....	674	<i>Marsdenia suaveolens</i> .....	968
<i>Narbonense</i> .....	190	<i>Maurandya Barclayana</i> .....	1381
<i>trigynum</i> .....	1193	<i>Maxillaria atropurpurea</i> ..	1877
<i>Liparia hirsuta</i> .....	33	<i>Barringtoniæ</i> ..	1824
<i>sphærica</i> .....	642	<i>Deppii</i> .....	1612
<i>Liparis elata</i> .....	1558	<i>galeata</i> .....	1645
<i>foliosa</i> .....	1097	<i>gracilis</i> .....	1837
<i>priochilus</i> .....	1751	<i>ochroleuca</i> .....	1904
<i>Lissanthe Daphnoides</i> .....	466	<i>punctata</i> .....	1914
<i>Listera cordata</i> .....	532	<i>racemosa</i> .....	1318
<i>Lithospermum purpuro-cœ-</i>		<i>Warreana</i> .....	1884
<i>ruleum</i> .....	1766	<i>Medicago arborea</i> .....	1379
<i>Lobelia cœrulea</i> .....	1206	<i>Melaleuca decussata</i> .....	1208
<i>corymbosa</i> .....	1298	<i>fulgens</i> .....	378
<i>fulgens</i> .....	1435	<i>hypericifolia</i> .....	199
<i>syphilitica</i> .....	955	<i>pulchella</i> .....	200
<i>Surinamensis</i> .....	749	<i>squamea</i> .....	412
<i>Logania floribunda</i> .....	1118	<i>squarrosa</i> .....	1130
<i>Lonicera flava</i> .....	338	<i>Thymifolia</i> .....	439
<i>flexuosa</i> .....	1037	<i>Melanthium junceum</i> .....	978
<i>Lophiola aurea</i> .....	1314	<i>Melasphærulea parviflora</i> ..	1444
<i>Lotus albidus</i> .....	1063	<i>Melastoma corymbosa</i> .....	984
<i>Luculia gratissima</i> .....	1919	<i>Malabathrica</i> ..	1932
<i>Lupinus lepidus</i> .....	1980	<i>Napalensis</i> .....	707
<i>Nootkatensis</i> .....	897	<i>villosa</i> .....	853
<i>Lychnis coronata</i> .....	1433	<i>Melissa Pyrenaica</i> .....	1853
<i>Suecica</i> .....	891	<i>Menziesia cœrulea</i> .....	164
<i>Lygodium scandens</i> .....	742	<i>polifolia nana</i> ..	1907
<i>Lysimachia longifolia</i> .....	1422	<i>Mesembryanthemum blan-</i>	
<i>Lythrum Græfferi</i> .....	1338	<i>dum</i> .....	599
<i>Macradenia lutescens</i> .....	1556	<i>bracteatum</i> ..	251
<i>Magnolia acuminata</i> .....	418	<i>coccineum</i> ..	1033
<i>Annonæfolia</i> .....	1072	<i>formosum</i> ..	1293
<i>cordata</i> .....	474	<i>gemmaeflorum</i> 1160	
<i>fuscata</i> .....	1073	<i>glomeratum</i> 1043	
<i>glaucæ</i> .....	215	<i>incomptum</i> 1311	
<i>grandiflora</i> .....	814	<i>incurvum</i> ..	1265
<i>pumila</i> .....	1555	<i>lacerum</i> ....	1279
<i>pyramidata</i> .....	1092	<i>linguseforme</i> 1307	
<i>yulan</i> .....	1187	<i>noctiflorum</i> -	495
<i>Mahernia diffusa</i> .....	187	<i>polyanthou</i> ..	1281
<i>grandiflora</i> .....	293	<i>Mespilus acuminata</i> .....	919
<i>incisa</i> .....	907	<i>Metrosideros hispida</i> .....	106
<i>Malaxis Liliifolia</i> .....	198	<i>semperflorens</i>	523
<i>Ophioglossoides</i> ..	1146	<i>speciosa</i> .....	285
<i>Malpighia aquifolia</i> .....	1079	<i>Mimosa sensitiva</i> .....	249
<i>angustifolia</i> .....	321	<i>Mimulus alatus</i> .....	410
<i>coccifera</i> .....	1317	<i>rivularis</i> .....	1575
<i>fucata</i> .....	1937	<i>roseus</i> .....	1976
<i>glabra</i> .....	533	<i>variegatus</i> .....	1872
<i>Malva Campanuloides</i> .....	1670	<i>Mirbelia dilatata</i> .....	1367



GENERAL INDEX.

	No.		No.
Mirbelia reticulata .....	1371	Ornithogalum Sternbergii..	1209
Mitchella repens .....	979	Orobus aureus .....	1955
Monarda punctata .....	1437	coccineus.....	883
purpurea .....	1396	Fischeri .....	1740
Monsonia speciosa .....	898	hirsutus .....	929
Musa coccinea .....	475	Lathyroides .....	698
paradisiaca.....	684	luteus .....	783
rosacea .....	615	sylvaticus .....	1181
Mussaenda glabra .....	1269	variegatus .....	1168
pubescens.....	451	Orontium aquaticum .....	402
Mutisia speciosa .....	1809	Orthrosanthus multiflorus..	1474
Myoporum parvifolium ....	837	Osbeckia Chinensis .....	665
Myosotis scorpioides .....	169	Osteospermum pisiferum ..	470
Myrsine retusa .....	409	Othonna Abrotanifolia ....	1698
Myrtus pimenta <i>latifolia</i> ..	178	flabellifolia .....	728
Nauclea adina .....	1284	Oxalis Bowiciana .....	1782
Neja gracilis.....	1814	Braziliensis .....	1962
Neottia elata.....	343	caprina .....	1109
picta .....	214	carnosa .....	1647
procera .....	776	cernua .....	1154
speciosa .....	838	cupræa .....	824
spiralis .....	931	Deppii .....	1500
Nepenthes distillatoria ....	1017	hirta .....	213
Nerine humilis .....	1674	Martiana .....	1523
undulata .....	1669	Mauritiana .....	1780
Nerium coccineum .....	894	papilionacea.....	1551
oleander <i>album</i> ....	700	tenella .....	1096
variegatum.....	666	tenuifolia .....	712
Nigritella angustifolia ....	1643	tetraphylla .....	790
Nitraria Schoberi .....	1395	trifurcata .....	1056
Octomeria graminifolia ....	1891	Oxylobium arborescens ....	163
Olea Europæa .....	456	cordifolium ....	937
fragrans .....	1786	Oxytropis campestris.....	1287
undulata .....	379	pilosa .....	544
Omphalea triandra .....	519	Pachypodium tuberosum ..	1676
Oncidium bifolium .....	1845	Pachysandra procumbens ..	910
Carthagenense ..	662	Pæonia mollis .....	1263
crispum .....	1854	<i>Moutan papaveracea</i> ..	547
divaricatum .....	1212	<i>rosea</i> .....	1035
flexuosum .....	424	<i>officinalis Sabini</i> ..	1075
Harrisoniæ .....	1917	Panax aculeatum.....	977
Papilio .....	1086	trifolium .....	405
pulchellum .....	1984	Pancratium Amancaes ....	1266
pumilum .....	1732	amœnum .....	286
Onoclea obtusilobata .....	1578	declinatum ....	558
Ononis fruticosa .....	1569	fragrans .....	834
rotundifolia .....	1496	Mexicanum ..	274
Onosma Echioides .....	1741	ovatum .....	510
rupestre.....	1880	pediale.....	809
Taurica .....	1405	rotatum .....	19
Oxylobium Pultenæ .....	1947	Papaver Alpinum.....	434
Ophiopogon Jaburan .....	1876	Pardanthus Chinensis ....	1874
spicatus .....	694	Passerina laxa .....	755
Ophrys Alpina.....	1188	spicata.....	311
Orchis fuscescens .....	1748	Passiflora alata.....	246
spectabilis .....	78	cœruleo-racemosa ..	573
Origanum Tournefortii ....	258	discolor .....	565
Ornithidium coccineum ....	301	fœtida .....	725
Ornithogalum Alliaceum ..	1818	Herbertiana .....	1364
aureum .....	1183	hirsuta .....	138
bifolium .....	1802	lunata .....	181
lacteum .....	1159	palmata .....	97
longibracteatum ..	1789	picturata.....	1050



GENERAL INDEX.

	No.		No.
<i>Passiflora princeps</i> .....	84	<i>Fimelea pauciflora</i> .....	179
<i>punctata</i> .....	101	<i>rosea</i> .....	88
<i>serratifolia</i> .....	195	<i>sylvestris</i> .....	1965
<i>Patersonia glabrata</i> .....	768	<i>Pinguicula Alpina</i> .....	309
<i>glauca</i> .....	1182	<i>grandiflora</i> .....	445
<i>Patrinia Scabiosæfolia</i> .....	1340	<i>vulgaris</i> .....	393
<i>Pavonia præmorsa</i> .....	371	<i>Piper coriaceum</i> .....	128
<i>Pelargonium ardens</i> .....	139	<i>discolor</i> .....	610
<i>Heracifolium</i> .....	437	<i>incanum</i> .....	503
<i>ignescens</i> .....	109	<i>maculosum</i> .....	861
<i>melananthos</i> .....	87	<i>pulchellum</i> .....	574
<i>Penæa marginata</i> .....	1770	<i>Pitcairnia staminea</i> .....	722
<i>Pentstemon angustifolium</i> ..	1420	<i>Pittosporum angustifolium</i>	1859
<i>atropurpureum</i> .....	1429	<i>coriaceum</i> ..	569
<i>gracilis</i> .....	1541	<i>revolutum</i> ..	506
<i>ovatum</i> .....	1626	<i>tomentosum</i> ..	1441
<i>procerum</i> .....	1616	<i>Plantago montana</i> .....	1581
<i>pulchellum</i> .....	1438	<i>Platylobium parviflorum</i> ..	1241
<i>Richardsonii</i> ..	1641	<i>triangulare</i> ..	1014
<i>Periploca Græca</i> .....	1389	<i>Plectranthus australis</i> .....	1185
<i>Persoonia flexifolia</i> .....	922	<i>Pleurothallis Lanceana</i> .....	1767
<i>hirsuta</i> .....	327	<i>prolifera</i> .....	1908
<i>lanceolata</i> .....	25	<i>racemiflora</i> ..	949
<i>latifolia</i> .....	1509	<i>saurocephalus</i> ..	1571
<i>spathulata</i> .....	1199	<i>Plumbago Capensis</i> .....	295
<i>Petiveria Alliacea</i> .....	148	<i>rhomboidea</i> .....	1536
<i>Petræa erecta</i> .....	1606	<i>Plumieria Mexicana</i> .....	1024
<i>volubilis</i> .....	736	<i>tuberculata</i> .....	681
<i>Petrocallis Pyrenaica</i> .....	635	<i>Podalyria Buxifolia</i> .....	1020
<i>Petrophyla trifida</i> .....	1883	<i>sericea</i> .....	1442
<i>Phaca Astragalina</i> .....	429	<i>Styracifolia</i> .....	649
<i>australis</i> .....	490	<i>Podolobium staurophyllum</i>	1177
<i>Phajus maculatus</i> .....	1803	<i>Polypodium pectinatum</i> ..	748
<i>Phlox aristata</i> .....	1731	<i>Polygala attenuata</i> .....	1000
<i>carnea</i> .....	711	<i>bracteolata</i> .....	1211
<i>crassifolia</i> .....	1596	<i>chamæbuxus</i> .....	593
<i>divaricata</i> .....	1418	<i>cordifolia</i> .....	1189
<i>nivalis</i> .....	780	<i>grandiflora</i> .....	1227
<i>pilosa</i> .....	1251	<i>Heisteria</i> .....	268
<i>procumbens</i> .....	1722	<i>humilis</i> .....	420
<i>pyramidalis</i> .....	342	<i>latifolia</i> .....	760
<i>Pholidota imbricata</i> .....	1934	<i>micrantha</i> .....	1773
<i>Phycella glauca</i> .....	1746	<i>seneka</i> .....	1380
<i>ignea</i> .....	1769	<i>speciosa</i> .....	621
<i>Phylla Buxifolia</i> .....	848	<i>stipulacea</i> .....	741
<i>plumosa</i> .....	253	<i>teretifolia</i> .....	294
<i>pubescens</i> .....	695	<i>Polygonum frutescens</i> .....	489
<i>Roëmarinifolia</i> .....	849	<i>viviparum</i> .....	1525
<i>spicata</i> .....	323	<i>Pomaderris elliptica</i> .....	756
<i>squarrosa</i> .....	36	<i>lanigera</i> .....	357
<i>Phyllanthus Emblia</i> .....	548	<i>Phylicifolia</i> .....	120
<i>Fraxinifolia</i> ..	839	<i>Pontederia lanceolata</i> .....	613
<i>Mimosoides</i> ..	721	<i>Ponthieva petiolata</i> .....	1190
<i>reticulata</i> .....	116	<i>Potentilla alba</i> .....	1534
<i>turbinata</i> .....	731	<i>Alpestris</i> .....	1499
<i>Phyteuma orbicularis</i> .....	122	<i>atrosanguinea</i> .....	786
<i>virgata</i> .....	667	<i>aurea</i> .....	77
<i>Pimelea decussata</i> .....	1283	<i>Calabra</i> .....	1757
<i>Diosmæfolia</i> .....	1703	<i>glabra</i> .....	914
<i>drupacea</i> .....	540	<i>Lupinaster</i> .....	654
<i>glauca</i> .....	1611	<i>minima</i> .....	480
<i>hispida</i> .....	1966	<i>Napalensis</i> .....	1031
<i>Linifolia</i> .....	1668	<i>nivea</i> .....	460



GENERAL INDEX.

	No.		No.
Potentilla pedata .....	579	Ranunculus amplexicaule..	1593
splendens .....	1323	Illyricus.....	1620
Weinmannii .....	706	montanus .....	1610
Pothos acaulis .....	483	Parnassifolius -	245
Cannæfolia .....	471	pedatus .....	351
Harrisii .....	1301	Rauvolfia nitida .....	339
microphylla.....	1673	Relhania pungens .....	1175
pentaphylla.....	567	Renealmia grandiflora .....	993
violacea .....	632	Rhamnus Alpinus .....	1077
Prescottia plantaginefolia..	990	Rhexia glomerata .....	334
Primula decora.....	1480	holosericea .....	236
farinosa .....	1640	Mariana .....	366
Helvetica .....	348	Virginica .....	840
inflata .....	1597	Rhipsalis cassutha .....	865
integrifolia .....	886	Mesembryanthemoides	1920
longiflora .....	542	Salicornoides .....	369
marginata .....	270	Rhododendron campanula-	
minima .....	315	tum.....	1944
Pallasii .....	1585	Catawbiense .....	1176
pusilla.....	1726	chamæcistus .....	1491
Scotica .....	652	Dauricum .....	605
Sinensis .....	916	var. ....	1446
<i>alba</i> .....	1926	<i>sempervirens</i>	1584
verticillata .....	1834	ferrugineum .....	65
villosa .....	182	hirsutum .....	479
Prinos glaber .....	450	Myrtifolium .....	908
Prockia Crucis.....	1933	Ribes alpinum .....	1486
Prostranthera violacea .....	1376	fragrans .....	1533
Prunus borealis .....	1598	lacustris .....	884
dasycarpa .....	1250	multiflorum .....	1331
depressa .....	1607	sanguineum .....	1487
Sibirica .....	1627	triflorum .....	1094
Psilotum triquetrum .....	1916	Robinia altagana .....	1064
Psoralea aphylla .....	221	jubata.....	522
bracteata .....	1559	Rodriguezia lanceolata .....	676
decumbens .....	282	Roella ciliata .....	1156
Pteris calomelanos .....	1689	Rondeletia hirta .....	350
palmata .....	1299	speciosa .....	1893
Peruviana .....	1665	Rosa Banksiæ <i>lutea</i> .....	1960
Pulmonaria Davurica .....	528	spinosissima <i>picta</i> .....	687
maritima .....	467	Roscoea purpurea .....	1404
mollis .....	1492	Rubus spectabilis .....	1602
Pultenæa biloba .....	550	Rosæfolius .....	158
candida .....	1236	Rudbeckia hirta .....	1386
daphnoides.....	1143	serotina.....	1539
mucronata .....	1711	triloba .....	817
paleacea .....	291	Ruellia anisophylla.....	1070
stipularis .....	1088	oblongifolia .....	1889
stricta .....	974	picta .....	1448
tenuifolia .....	1057	Sabiniana .....	1712
Punica nana .....	988	Rulingia pannosa .....	491
Pyrethrum saxatile .....	126	Ruta albiflora .....	1352
Pyrola chlorantha .....	1542	Salpiglossis integrifolia .....	1978
maculata .....	708	picta.....	1652
rotundifolia .....	564	Salvia amœna .....	377
umbellata .....	463	Chamædryoides .....	576
Pyrus Japonica .....	1594	fulgens .....	1910
<i>alba</i> .....	541	Grahami .....	1798
Pollveria .....	1009	splendens .....	1089
Salicifolia .....	1120	Samolus littoralis .....	435
spectabilis .....	1729	Sanguinaria Canadensis....	1840
Quassia amara .....	172	Sarracenia flava .....	1957
Rafnia triflora .....	611	purpurea .....	308



GENERAL INDEX.

	No.		No.
Tacca pinnatifida .....	692	Vaccinium vitis-idaea <i>major</i>	616
Talinum teretifolium .....	819	<i>minor</i>	1023
Tarhonanthus camphoratus .....	382	Valantia Taurica .....	1018
Tecoma capensis .....	1672	Valeriana montana .....	317
Templetonia glauca .....	644	Vanda rostrata .....	1008
retusa .....	526	Vanilla planifolia .....	733
Teucrium orientale .....	1871	Veltheimia viridifolia .....	1245
Pyrenaicum .....	1387	Verbascum Phœniceum .....	637
Thalictrum petaloideum ..	891	Verbena Melindris .....	1514
Thea Bohea .....	226	pulchella .....	1550
viridis .....	227	Vernonia axillæflora .....	1690
<i>latifolia</i> .....	1828	flexuosa .....	1680
Theobroma Cacao .....	545	Veronica Caucasica .....	1369
Thermopsis Fabacea .....	1856	Chamædrys .....	53
Laburnifolia ..	1095	incisa .....	1897
Thuja articulata .....	844	orientalis .....	419
Thunbergia alata .....	1045	perfoliata .....	781
angulata .....	1044	saxatilis .....	704
capensis .....	1529	Taurica .....	911
coccinea .....	1195	Teucrium .....	425
fragrans .....	1913	Viburnum Lantanoides .....	1570
grandiflora .....	324	obovatum .....	1476
Thymus Azoricus .....	1530	oxycoccos .....	1123
Patavinus .....	499	rugosum .....	859
Thysanotus junceus .....	1936	Vieusseuxia glaucopsis .....	1744
Tigridia Pavonia .....	1424	Villarsia Chilensis .....	1994
Tillandsia amœna .....	76	Viola debilis .....	1378
anceps .....	771	dentata .....	1485
Torenia scabra .....	1990	flabellifolia .....	777
Tradescantia alata .....	1513	Hederacea .....	1133
crassula .....	1560	lanceolata .....	211
cristata .....	639	pedata .....	536
fuscata .....	374	pubescens .....	1249
<i>latifolia</i> .....	1300	Rupii .....	686
rosea .....	370	sagittata .....	1471
Trichonema cruciatum .....	1370	Vismea glabra .....	1752
Trientalis Europæa .....	105	Uvularia puberula .....	1260
Trifolium fimbriatum .....	1421	sessilifolia .....	1262
uniflorum .....	1882	Waldsteinia Geoides .....	492
Triglochin bulbosa .....	1151	Watsonia compacta .....	1577
Trigonella Ruthenica .....	1391	Westringia angustifolia .....	1890
Trillium erectum .....	1838	Witheringia montana .....	1292
<i>album</i> ..	1850	purpurea .....	1892
erythrocarpum ..	1232	Witsenia corymbosa .....	254
grandiflorum .....	1349	Wrightia pubescens .....	1929
nervosum .....	1860	Xerotis rigida .....	798
sessile .....	875	Xylophylla angustifolia .....	577
Tristania Neriifolia .....	157	elongata .....	1091
Tritoma pumila .....	444	Xylosteum campaniflorum	1361
Tritonia odorata .....	1820	Xyris altissima .....	1900
tenuiflora .....	1078	operculata .....	205
Trizeuxis falcata .....	975	Zamia debilis .....	155
Trollius Americanus .....	56	pygmæa .....	113
Tropæolum pentaphyllum ..	1989	Zephyranthes Atamasco ..	1899
Tupistra squalida .....	515	candida .....	1419
Tussilago alpina .....	1618	Zieria lanceolata .....	878
Vaccinium amœnum .....	774	Zygopetalum crinitum .....	1687
arboreum .....	1885	Mackaii .....	1664
Buxifolium .....	648	maxillare .....	1776
ovatum .....	1605	stenochilum ..	1923

THE END.

W. WILSON, PRINTER, 57, SKINNER-STREET, LONDON.